arXiv:1207.6162v1 [math-ph] 26 Jul 2012

Cyclic structures of Cliffordian supergroups and particle representations of $\mathbf{Spin}_+(1,3)$

V. V. Varlamov

Department of Mathematics, Siberian State Industrial University,
Kirova 42, Novokuznetsk 654007, Russia

Abstract

Supergroups are defined in the framework of $\mathbb{Z}_2$-graded Clifford algebras over the fields of real and complex numbers, respectively. It is shown that cyclic structures of complex and real supergroups are defined by Brauer-Wall groups related with the modulo 2 and modulo 8 periodicities of the complex and real Clifford algebras. Particle (fermionic and bosonic) representations of a universal covering (spinor group $\mathbf{Spin}_+(1,3)$) of the proper orthochronous Lorentz group are constructed via the Clifford algebra formalism. Complex and real supergroups are defined on the representation system of $\mathbf{Spin}_+(1,3)$. It is shown that a cyclic (modulo 2) structure of the complex supergroup is equivalent to a supersymmetric action, that is, it converts fermionic representations into bosonic representations and vice versa. The cyclic action of the real supergroup leads to a much more high-graded symmetry related with the modulo 8 periodicity of the real Clifford algebras. This symmetry acts on the system of real representations of $\mathbf{Spin}_+(1,3)$.

Keywords: Clifford algebras, supergroups, spinor representations
MSC 2010: **15A66, 15A90, 20645**
PACS numbers: **02.10.Tq, 11.30.Er, 11.30.Cp**

1 Introduction

It is well known that Clifford algebras (systems of hypercomplex numbers) have a broad application in many areas of mathematical and theoretical physics. From historical point of view, Clifford algebras have essentially geometric origin [1], because they are the synthesis of Hamilton quaternion calculus [2] and Grassmann *Ausdehnungslehre* [3], and by this reason they called by Clifford as *geometric algebras* [4]. Further, Lipschitz [5] showed that Clifford algebras are related closely with the study of rotation groups of multidimensional spaces. After fundamental works of Cartan [6], Witt [7] and Chevalley [8] the Clifford algebra theory takes its modern form [9, 10, 11]. It is known that Clifford algebras inherit $\mathbb{Z}_2$-graded structure from the Grassmann algebras. This fact allows one to define *Cliffordian supergroups* using a classical theory of formal Lie groups [12, 13]. Along with the $\mathbb{Z}_2$-graded structure Clifford algebras possess modulo 2 and modulo 8 periodicities over the fields of complex and real numbers, respectively (see [14, 15, 16]). These periodic relations are described by Brauer-Wall groups [17, 18]. The cyclic (modulo 2 and 8) structure is the most essential property of the Cliffordian supergroups.

In the present work we consider the field $\boldsymbol{\psi}(\boldsymbol{\alpha}) = \langle x, \mathfrak{g}\,|\boldsymbol{\psi}\rangle$ on the representation spaces of a spinor group $\mathbf{Spin}_+(1,3)$, where $x \in T_4$ and $\mathfrak{g} \in \mathbf{Spin}_+(1,3)$ (T_4 is a translation subgroup of the Poincaré group $\mathcal{P}$). At this point, four parameters x^μ correspond to position of the point-like object, whereas remaining six parameters $\mathbf{z} \in \mathbf{Spin}_+(1,3)$ define orientation in quantum description of orientable (extended) object [19, 20]. In general, the field $\boldsymbol{\psi}(\boldsymbol{\alpha})$ is defined on the homogeneous space $\mathcal{M}_{10} = \mathbb{R}^{1,3} \times \mathfrak{L}_6$ (a group manifold of $\mathcal{P}$), where $\mathbb{R}^{1,3}$ is the Minkowski spacetime and $\mathfrak{L}_6$ is a group manifold of $\mathrm{SO}_0(1,3)$. On the other hand, the space $\mathcal{M}_{10} = \mathbb{R}^{1,3} \times \mathfrak{L}_6$ can be understood as a fiber bundle, where a bundle base is $\mathbb{R}^{1,3}$. All intrinsic properties of the field $\boldsymbol{\psi}(\boldsymbol{\alpha})$ are described within the group $\mathbf{Spin}_+(1,3)$ and its representations.

The paper is organized as follows. A short introduction to the Clifford algebra theory is given in section 2. $\mathbb{Z}_2$-graded Clifford algebras, Brauer-Wall supergroups and Trautman diagrams (spinorial clocks) are considered in section 3 over the fields of real and complex numbers. Complex and real representations of the field $\boldsymbol{\psi}(\boldsymbol{\alpha}) = \langle x, \mathfrak{g} \,|\boldsymbol{\psi}\rangle$ of different types are constructed in section 4 within representations of $\mathbf{Spin}_+(1,3)$. A relationship between tensor products of the Clifford algebras and a Gel'fand-Naimark representation basis of the proper orthochronous Lorentz group is established. Complex and real supergroups are defined on the representation systems of $\mathbf{Spin}_+(1,3)$ in section 5.

2 Algebraic preliminaries

In this section we will consider some basic facts concerning Clifford algebras. Let $\mathbb{F}$ be a field of characteristic 0 ($\mathbb{F} = \mathbb{R}$, $\mathbb{F} = \mathbb{C}$), where $\mathbb{R}$ and $\mathbb{C}$ are the fields of real and complex numbers, respectively. A Clifford algebra $C\!\ell$ over a field $\mathbb{F}$ is an algebra with 2^n basis elements: $\mathbf{e}_0$ (unit of the algebra) $\mathbf{e}_1, \mathbf{e}_2, \ldots, \mathbf{e}_n$ and products of the one-index elements $\mathbf{e}_{i_1 i_2 \ldots i_k} = \mathbf{e}_{i_1}\mathbf{e}_{i_2}\ldots\mathbf{e}_{i_k}$. Over the field $\mathbb{F} = \mathbb{R}$ the Clifford algebra is denoted as $C\!\ell_{p,q}$, where the indices p and q correspond to the indices of the quadratic form

$$Q = x_1^2 + \ldots + x_p^2 - \ldots - x_{p+q}^2$$

of a vector space V associated with $C\!\ell_{p,q}$. The multiplication law of $C\!\ell_{p,q}$ is defined by a following rule:

$$\mathbf{e}_i^2 = \sigma(p-i)\mathbf{e}_0, \quad \mathbf{e}_i\mathbf{e}_j = -\mathbf{e}_j\mathbf{e}_i, \tag{1}$$

where

$$\sigma(n) = \begin{cases} -1 & \text{if } n \leq 0, \\ +1 & \text{if } n > 0. \end{cases} \tag{2}$$

The square of a volume element $\omega = \mathbf{e}_{12\ldots n}$ $(n = p+q)$ plays an important role in the theory of Clifford algebras,

$$\omega^2 = \begin{cases} -1 & \text{if } p-q \equiv 2,3,6,7 \pmod 8, \\ +1 & \text{if } p-q \equiv 0,1,4,5 \pmod 8. \end{cases} \tag{3}$$

A center $\mathbf{Z}_{p,q}$ of the algebra $C\!\ell_{p,q}$ consists of the unit $\mathbf{e}_0$ and the volume element ω. The element $\omega = \mathbf{e}_{12\ldots n}$ belongs to a center when n is odd. Indeed,

$$\begin{aligned} \mathbf{e}_{12\ldots n}\mathbf{e}_i &= (-1)^{n-i}\sigma(q-i)\mathbf{e}_{12\ldots i-1 i+1\ldots n}, \\ \mathbf{e}_i\mathbf{e}_{12\ldots n} &= (-1)^{i-1}\sigma(q-i)\mathbf{e}_{12\ldots i-1 i+1\ldots n}, \end{aligned}$$

therefore, $\omega \in \mathbf{Z}_{p,q}$ if and only if $n - i \equiv i - 1 \pmod 2$, that is, n is odd. Further, using (3) we obtain

$$\mathbf{Z}_{p,q} = \begin{cases} 1 & \text{if } p-q \equiv 0,2,4,6 \pmod 8, \\ 1, \omega & \text{if } p-q \equiv 1,3,5,7 \pmod 8. \end{cases} \tag{4}$$

An arbitrary element $\mathcal{A}$ of the algebra $C\!\ell_{p,q}$ is represented by a following formal polynomial

$$\mathcal{A}=a^0\mathbf{e}_0+\sum_{i=1}^n a^i\mathbf{e}_i+\sum_{i=1}^n\sum_{j=1}^n a^{ij}\mathbf{e}_{ij}+\ldots+\sum_{i_1=1}^n\cdots\sum_{i_k=1}^n a^{i_1\ldots i_k}\mathbf{e}_{i_1\ldots i_k}+$$
$$+\ldots+a^{12\ldots n}\mathbf{e}_{12\ldots n}=\sum_{k=0}^n a^{i_1i_2\ldots i_k}\mathbf{e}_{i_1i_2\ldots i_k}.$$

In Clifford algebra $C\!\ell$ there exist four fundamental automorphisms.

1) **Identity**: An automorphism $\mathcal{A}\rightarrow\mathcal{A}$ and $\mathbf{e}_i\rightarrow\mathbf{e}_i$.
This automorphism is an identical automorphism of the algebra $C\!\ell$. $\mathcal{A}$ is an arbitrary element of $C\!\ell$.

2) **Involution**: An automorphism $\mathcal{A}\rightarrow\mathcal{A}^\star$ and $\mathbf{e}_i\rightarrow-\mathbf{e}_i$.
In more details, for an arbitrary element $\mathcal{A}\in C\!\ell$ there exists a decomposition $\mathcal{A}=\mathcal{A}'+\mathcal{A}''$, where $\mathcal{A}'$ is an element consisting of homogeneous odd elements, and $\mathcal{A}''$ is an element consisting of homogeneous even elements, respectively. Then the automorphism $\mathcal{A}\rightarrow\mathcal{A}^\star$ is such that the element $\mathcal{A}''$ is not changed, and the element $\mathcal{A}'$ changes sign: $\mathcal{A}^\star=-\mathcal{A}'+\mathcal{A}''$. If $\mathcal{A}$ is a homogeneous element, then

$$\mathcal{A}^\star=(-1)^k\mathcal{A}, \tag{5}$$

where k is a degree of the element. It is easy to see that the automorphism $\mathcal{A}\rightarrow\mathcal{A}^\star$ may be expressed via the volume element $\omega=\mathbf{e}_{12\ldots p+q}$:

$$\mathcal{A}^\star=\omega\mathcal{A}\omega^{-1}, \tag{6}$$

where $\omega^{-1}=(-1)^{\frac{(p+q)(p+q-1)}{2}}\omega$. When k is odd, the basis elements $\mathbf{e}_{i_1i_2\ldots i_k}$ the sign changes, and when k is even, the sign is not changed.

3) **Reversion**: An antiautomorphism $\mathcal{A}\rightarrow\widetilde{\mathcal{A}}$ and $\mathbf{e}_i\rightarrow\mathbf{e}_i$.
The antiautomorphism $\mathcal{A}\rightarrow\widetilde{\mathcal{A}}$ is a reversion of the element $\mathcal{A}$, that is the substitution of each basis element $\mathbf{e}_{i_1i_2\ldots i_k}\in\mathcal{A}$ by the element $\mathbf{e}_{i_ki_{k-1}\ldots i_1}$:

$$\mathbf{e}_{i_ki_{k-1}\ldots i_1}=(-1)^{\frac{k(k-1)}{2}}\mathbf{e}_{i_1i_2\ldots i_k}.$$

Therefore, for any $\mathcal{A}\in C\!\ell_{p,q}$ we have

$$\widetilde{\mathcal{A}}=(-1)^{\frac{k(k-1)}{2}}\mathcal{A}. \tag{7}$$

4) **Conjugation**: An antiautomorphism $\mathcal{A}\rightarrow\widetilde{\mathcal{A}^\star}$ and $\mathbf{e}_i\rightarrow-\mathbf{e}_i$.
This antiautomorphism is a composition of the antiautomorphism $\mathcal{A}\rightarrow\widetilde{\mathcal{A}}$ with the automorphism $\mathcal{A}\rightarrow\mathcal{A}^\star$. In the case of a homogeneous element from the formulae (5) and (7), it follows

$$\widetilde{\mathcal{A}^\star}=(-1)^{\frac{k(k+1)}{2}}\mathcal{A}. \tag{8}$$

The all Clifford algebras $C\!\ell_{p,q}$ over the field $\mathbb{F}=\mathbb{R}$ are divided into eight different types with a following division ring structure.

I. Central simple algebras.

1) Two types $p-q\equiv 0,2 \pmod 8$ with a division ring $\mathbb{K}\simeq\mathbb{R}$.

2) Two types $p-q\equiv 3,7 \pmod 8$ with a division ring $\mathbb{K}\simeq\mathbb{C}$.

p	0	1	2	3	4	5	6	7	$\ldots$
q									
0	$\mathbb{R}$	${}^2\mathbb{R}$	$\mathbb{R}(2)$	$\mathbb{C}(2)$	$\mathbb{H}(2)$	${}^2\mathbb{H}(2)$	$\mathbb{H}(4)$	$\mathbb{C}(8)$	$\ldots$
1	$\mathbb{C}$	$\mathbb{R}(2)$	${}^2\mathbb{R}(2)$	$\mathbb{R}(4)$	$\mathbb{C}(4)$	$\mathbb{H}(4)$	${}^2\mathbb{H}(4)$	$\mathbb{H}(8)$	$\ldots$
2	$\mathbb{H}$	$\mathbb{C}(2)$	$\mathbb{R}(4)$	${}^2\mathbb{R}(4)$	$\mathbb{R}(8)$	$\mathbb{C}(8)$	$\mathbb{H}(8)$	${}^2\mathbb{H}(8)$	$\ldots$
3	${}^2\mathbb{H}$	$\mathbb{H}(2)$	$\mathbb{C}(4)$	$\mathbb{R}(8)$	${}^2\mathbb{R}(8)$	$\mathbb{R}(16)$	$\mathbb{C}(16)$	$\mathbb{H}(16)$	$\ldots$
4	$\mathbb{H}(2)$	${}^2\mathbb{H}(2)$	$\mathbb{H}(4)$	$\mathbb{C}(8)$	$\mathbb{R}(16)$	${}^2\mathbb{R}(16)$	$\mathbb{R}(32)$	$\mathbb{C}(32)$	$\ldots$
5	$\mathbb{C}(4)$	$\mathbb{H}(4)$	${}^2\mathbb{H}(4)$	$\mathbb{H}(8)$	$\mathbb{C}(16)$	$\mathbb{R}(32)$	${}^2\mathbb{R}(32)$	$\mathbb{R}(64)$	$\ldots$
6	$\mathbb{R}(8)$	$\mathbb{C}(8)$	$\mathbb{H}(8)$	${}^2\mathbb{H}(8)$	$\mathbb{H}(16)$	$\mathbb{C}(32)$	$\mathbb{R}(64)$	${}^2\mathbb{R}(64)$	$\ldots$
7	${}^2\mathbb{R}(8)$	$\mathbb{R}(16)$	$\mathbb{C}(16)$	$\mathbb{H}(16)$	${}^2\mathbb{H}(16)$	$\mathbb{H}(32)$	$\mathbb{C}(64)$	$\mathbb{R}(128)$	$\ldots$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	

Tab. 1: Distribution of the real Clifford algebras.

3) Two types $p - q \equiv 4, 6 \pmod 8$ with a division ring $\mathbb{K} \simeq \mathbb{H}$.

II. Semi-simple algebras.

4) The type $p - q \equiv 1 \pmod 8$ with a double division ring $\mathbb{K} \simeq \mathbb{R} \oplus \mathbb{R}$.

5) The type $p - q \equiv 5 \pmod 8$ with a double quaternionic division ring $\mathbb{K} \simeq \mathbb{H} \oplus \mathbb{H}$.

The Table 1 (Budinich-Trautman Periodic Table [16]) explicitly shows a distribution of the real Clifford algebras in dependence on the division ring structure, here ${}^2\mathbb{R}(n) = \mathbb{R}(n) \oplus \mathbb{R}(n)$ and ${}^2\mathbb{H}(n) = \mathbb{H}(n) \oplus \mathbb{H}(n)$.

Over the field $\mathbb{F} = \mathbb{C}$ there is an isomorphism $\mathbb{C}_n \simeq \mathrm{Mat}_{2^{n/2}}(\mathbb{C})$ and there are two different types of complex Clifford algebras $\mathbb{C}_n$: $n \equiv 0 \pmod 2$ and $n \equiv 1 \pmod 2$.

When $\mathcal{C}\ell_{p,q}$ is simple, then the map

$$\mathcal{C}\ell_{p,q} \xrightarrow{\gamma} \mathrm{End}_{\mathbb{K}}(\mathbb{S}), \quad u \longrightarrow \gamma(u), \quad \gamma(u)\psi = u\psi \tag{9}$$

gives an irreducible and faithful representation of $\mathcal{C}\ell_{p,q}$ in the spinspace $\mathbb{S}_{2^m}(\mathbb{K}) \simeq I_{p,q} = \mathcal{C}\ell_{p,q} f$, where $\psi \in \mathbb{S}_{2^m}$, $m = \frac{p+q}{2}$, $I_{p,q}$ is a minimal left ideal of $\mathcal{C}\ell_{p,q}$.

On the other hand, when $\mathcal{C}\ell_{p,q}$ is semi-simple, then the map

$$\mathcal{C}\ell_{p,q} \xrightarrow{\gamma} \mathrm{End}_{\mathbb{K}\oplus\hat{\mathbb{K}}}(\mathbb{S} \oplus \hat{\mathbb{S}}), \quad u \longrightarrow \gamma(u), \quad \gamma(u)\psi = u\psi \tag{10}$$

gives a faithful but reducible representation of $\mathcal{C}\ell_{p,q}$ in the double spinspace $\mathbb{S} \oplus \hat{\mathbb{S}}$, where $\hat{\mathbb{S}} = \{\hat{\psi} | \psi \in \mathbb{S}\}$. In this case, the ideal $\mathbb{S} \oplus \hat{\mathbb{S}}$ possesses a right $\mathbb{K} \oplus \hat{\mathbb{K}}$-linear structure, $\hat{\mathbb{K}} = \{\hat{\lambda} | \lambda \in \mathbb{K}\}$, and $\mathbb{K} \oplus \hat{\mathbb{K}}$ is isomorphic to the double division ring $\mathbb{R} \oplus \mathbb{R}$ when $p - q \equiv 1 \pmod 8$ or to $\mathbb{H} \oplus \mathbb{H}$ when $p - q \equiv 5 \pmod 8$. The map γ in (9) and (10) defines the so called *left-regular* spinor representation of $\mathcal{C}\ell(Q)$ in $\mathbb{S}$ and $\mathbb{S} \oplus \hat{\mathbb{S}}$, respectively. Furthermore, γ is *faithful* which means that γ is an algebra monomorphism. In (9), γ is *irreducible* which means that $\mathbb{S}$ possesses no proper (that is, $\neq 0$, $\mathbb{S}$) invariant subspaces under the left action of $\gamma(u)$, $u \in \mathcal{C}\ell_{p,q}$. Representation γ in (10) is therefore *reducible* since $\{(\psi, 0) | \psi \in \mathbb{S}\}$ and $\{(0, \hat{\psi}) | \hat{\psi} \in \hat{\mathbb{S}}\}$ are two proper subspaces of $\mathbb{S} \oplus \hat{\mathbb{S}}$ invariant under $\gamma(u)$ (see [11, 9, 10]).

3 $\mathbb{Z}_2$-graded Clifford algebras and Brauer-Wall groups

The algebra $\mathcal{C}\!\ell$ is naturally $\mathbb{Z}_2$-graded. Let $\mathcal{C}\!\ell^+$ (correspondingly $\mathcal{C}\!\ell^-$) be a set consisting of all even (correspondingly odd) elements of the algebra $\mathcal{C}\!\ell$. The set $\mathcal{C}\!\ell^+$ is a subalgebra of $\mathcal{C}\!\ell$. It is obvious that $\mathcal{C}\!\ell = \mathcal{C}\!\ell^+ \oplus \mathcal{C}\!\ell^-$, and also $\mathcal{C}\!\ell^+\mathcal{C}\!\ell^+ \subset \mathcal{C}\!\ell^+$, $\mathcal{C}\!\ell^+\mathcal{C}\!\ell^- \subset \mathcal{C}\!\ell^-$, $\mathcal{C}\!\ell^-\mathcal{C}\!\ell^+ \subset \mathcal{C}\!\ell^-$, $\mathcal{C}\!\ell^-\mathcal{C}\!\ell^- \subset \mathcal{C}\!\ell^+$. A degree $\deg a$ of the even (correspondingly odd) element $a \in \mathcal{C}\!\ell$ is equal to 0 (correspondingly 1). Let $\mathfrak{A}$ and $\mathfrak{B}$ be the two associative $\mathbb{Z}_2$-graded algebras over the field $\mathbb{F}$; then a multiplication of homogeneous elements $\mathfrak{a}' \in \mathfrak{A}$ and $\mathfrak{b} \in \mathfrak{B}$ in a graded tensor product $\mathfrak{A}\hat{\otimes}\mathfrak{B}$ is defined as follows: $(\mathfrak{a} \otimes \mathfrak{b})(\mathfrak{a}' \otimes \mathfrak{b}') = (-1)^{\deg \mathfrak{b} \deg \mathfrak{a}'}\mathfrak{a}\mathfrak{a}' \otimes \mathfrak{b}\mathfrak{b}'$.

Theorem 1 (Chevalley [21])**.** *Let V and V' are vector spaces over the field $\mathbb{F}$ and let Q and Q' are quadratic forms for V and V'. Then a Clifford algebra $\mathcal{C}\!\ell(V \oplus V', Q \oplus Q')$ is naturally isomorphic to $\mathcal{C}\!\ell(V,Q)\hat{\otimes}\mathcal{C}\!\ell(V',Q')$.*

Let $\mathcal{C}\!\ell(V,Q)$ be the Clifford algebra over the field $\mathbb{F} = \mathbb{R}$, where V is a vector space endowed with quadratic form $Q = x_1^2 + \ldots + x_p^2 - \ldots - x_{p+q}^2$. If $p+q$ is even and $\omega^2 = 1$, then $\mathcal{C}\!\ell(V,Q)$ is called *positive* and correspondingly *negative* if $\omega^2 = -1$, that is, $\mathcal{C}\!\ell_{p,q} > 0$ if $p - q \equiv 0, 4 \pmod 8$ and $\mathcal{C}\!\ell_{p,q} < 0$ if $p - q \equiv 2, 6 \pmod 8$.

Theorem 2 (Karoubi [22, Prop. 3.16])**.** *1) If $\mathcal{C}\!\ell(V,Q) > 0$ and $\dim V$ is even, then*

$$\mathcal{C}\!\ell(V \oplus V', Q \oplus Q') \simeq \mathcal{C}\!\ell(V,Q) \otimes \mathcal{C}\!\ell(V',Q').$$

2) If $\mathcal{C}\!\ell(V,Q) < 0$ and $\dim V$ is even, then

$$\mathcal{C}\!\ell(V \oplus V', Q \oplus Q') \simeq \mathcal{C}\!\ell(V,Q) \otimes \mathcal{C}\!\ell(V',-Q').$$

Over the field $\mathbb{F} = \mathbb{C}$ the Clifford algebra is always positive (if $\omega^2 = \mathbf{e}_{12\ldots p+q}^2 = -1$, then we can suppose $\omega = i\mathbf{e}_{12\ldots p+q}$). Thus, using Karoubi Theorem, we find that

$$\underbrace{\mathbb{C}_2 \otimes \cdots \otimes \mathbb{C}_2}_{m\,\text{times}} \simeq \mathbb{C}_{2m}. \tag{11}$$

Therefore, the tensor product in (11) is isomorphic to $\mathbb{C}_{2m}$. For example, there are two different factorizations $\mathcal{C}\!\ell_{1,1} \otimes \mathcal{C}\!\ell_{0,2}$ and $\mathcal{C}\!\ell_{1,1} \otimes \mathcal{C}\!\ell_{2,0}$ for the spacetime algebra $\mathcal{C}\!\ell_{1,3}$ and Majorana algebra $\mathcal{C}\!\ell_{3,1}$.

3.1 Supergroups

Let us consider graded tensor products $\mathfrak{A}_1\hat{\otimes}\mathfrak{A}_2$ and $\mathfrak{A}_1\hat{\otimes}\mathfrak{A}_2\hat{\otimes}\mathfrak{A}_3$ over the field $\mathbb{F}$ of characteristics 0, where $\mathfrak{A} = \mathcal{C}\!\ell_{p,q}$ over the field $\mathbb{F} = \mathbb{R}$ and $\mathfrak{A} = \mathbb{C}_n$ over $\mathbb{F} = \mathbb{C}$. These products possess the same properties as the algebra $\mathfrak{A}$. Let $\mathbf{e}_i'$ $(i = 1, \ldots, n_1)$, $\mathbf{e}_i''$ $(i = 1, \ldots, n_2)$, $\mathbf{e}_i'''$ $(i = 1, \ldots, n_3)$ be the units of the algebras $\mathfrak{A}_i$. Then the units of the algebras $\mathfrak{A}_1\hat{\otimes}\mathfrak{A}_2$ and $\mathfrak{A}_1\hat{\otimes}\mathfrak{A}_2\hat{\otimes}\mathfrak{A}_3$ are the sets $\mathbf{e}_i'$, $\mathbf{e}_j''$ and $\mathbf{e}_i'$, $\mathbf{e}_j''$, $\mathbf{e}_k'''$, respectively. General elements of the algebras $\mathfrak{A}_1\hat{\otimes}\mathfrak{A}_2$ and $\mathfrak{A}_1\hat{\otimes}\mathfrak{A}_2\hat{\otimes}\mathfrak{A}_3$ are formal polynomials from $n_1 + n_2$ and $n_1 + n_2 + n_3$ units. Dimensionalities of $\mathfrak{A}_1\hat{\otimes}\mathfrak{A}_2$ and $\mathfrak{A}_1\hat{\otimes}\mathfrak{A}_2\hat{\otimes}\mathfrak{A}_3$ are equal to $2^{n_1+n_2}$ and $2^{n_1+n_2+n_3}$, respectively.

Let d be a mapping of $\mathfrak{A}\hat{\otimes}\mathfrak{A}$ into $\mathfrak{A}$ (d defines a multiplication operation in the algebra $\mathfrak{A}$: $d_{a_1\otimes a_2} = a_1a_2$). Further, let us suppose $\eta(k) = k\mathbf{e}_0$, where $\mathbf{e}_0$ is the unit of the algebra $\mathfrak{A}$, $k \in \mathbb{F}$, and $\varepsilon_a = a(0)$. The mapping η defines a natural embedding of the field $\mathbb{F}$ into $\mathfrak{A}$, ε is an homomorphism of the algebra $\mathfrak{A}$ into the field $\mathbb{F}$.

Definition 1. *The algebra $\mathfrak{A}$ is called a supergroup space if there exist the homomorphism γ : $\mathfrak{A} \to \mathfrak{A}\hat{\otimes}\mathfrak{A}$ and an isomorphism $\bigcirc$: $\mathfrak{A} \to \mathfrak{A}$ with the following commutative diagrams:*

$$\begin{tikzcd}
 & \mathfrak{A}\hat{\otimes}\mathfrak{A} \arrow[dr, "I\otimes\gamma"] & \\
\mathfrak{A} \arrow[ur, "\gamma"] \arrow[dr, "\gamma"'] & & \mathfrak{A}\hat{\otimes}\mathfrak{A}\hat{\otimes}\mathfrak{A} \\
 & \mathfrak{A}\hat{\otimes}\mathfrak{A} \arrow[ur, "\gamma\otimes I"'] &
\end{tikzcd} \tag{12}$$

$$\begin{tikzcd}
 & \mathfrak{A}\hat{\otimes}\mathfrak{A} \arrow[dr, "I\otimes\varepsilon"] & \\
\mathfrak{A} \arrow[ur, "\gamma"] \arrow[rr, "I"'] & & \mathfrak{A}
\end{tikzcd} \tag{13}$$

$$\begin{tikzcd}
\mathfrak{A}\hat{\otimes}\mathfrak{A} \arrow[rr, "I\otimes\bigcirc"] & & \mathfrak{A}\hat{\otimes}\mathfrak{A} \arrow[d, "d"] \\
\mathfrak{A} \arrow[u, "\gamma"] \arrow[r, "\varepsilon"'] & \mathbb{F} \arrow[r, "\eta"'] & \mathfrak{A}
\end{tikzcd} \tag{14}$$

Definition 2. *The totality $G(\mathfrak{A}, \gamma, \bigcirc)$ is called a Cliffordian supergroup.*

The homomorphism γ is defined as follows. Let

$$\begin{aligned} j: &\quad V \longrightarrow \mathcal{C}\ell(V, Q), \\ j': &\quad V' \longrightarrow \mathcal{C}\ell(V', Q') \end{aligned}$$

be canonical mappings (linear mappings of the vector spaces into the algebras). Let us assume that V and V' are mutually orthogonal, then $V \oplus V'$. Let

$$j'': \; V \oplus V' \longrightarrow \mathcal{C}\ell(V, Q)\hat{\otimes}\mathcal{C}\ell(V', Q')$$

be a canonical mapping of the direct sum $V \oplus V'$ into the graded tensor product of the Clifford algebras defined by the formula $j''(v, v') = j(v) \otimes 1 + 1 \otimes j(v')$, where v and v' are vectors of the spaces V and V', respectively. Hence it immediately follows that $(j''(v, v'))^2 = [(j(v))^2 + (j'(v'))^2] \cdot 1 = [Q(v) + Q'(v')] \cdot 1$, since v and v' are orthogonal. Thus, the mapping j'' induces an homomorphism

$$\psi: \; \mathcal{C}\ell(V \oplus V', Q \oplus Q') \longrightarrow \mathcal{C}\ell(V, Q)\hat{\otimes}\mathcal{C}\ell(V', Q').$$

Let

$$\gamma: \quad \mathcal{C}\ell(V, Q) \longrightarrow \mathcal{C}\ell(V \oplus V', Q \oplus Q'), \tag{15}$$

$$\gamma': \quad \mathcal{C}\ell(V', Q') \longrightarrow \mathcal{C}\ell(V \oplus V', Q \oplus Q') \tag{16}$$

be homomorphisms induced by embeddings of V and V' into $V \oplus V'$. Since V and V' are orthogonal in $V \oplus V'$, then there exists an equality $\gamma(x)\gamma'(x') = (-1)^{\deg x \deg x'}\gamma'(x')\gamma(x)$, where x and x' are homogeneous elements of the algebras $\mathcal{C}\ell(V, Q)$ and $\mathcal{C}\ell(V', Q')$. Therefore, γ and γ' define an homomorphism

$$\theta: \; \mathcal{C}\ell(V, Q)\hat{\otimes}\mathcal{C}\ell(V', Q') \longrightarrow \mathcal{C}\ell(V \oplus V', Q \oplus Q')$$

by the formula $\theta(x\otimes x') = \gamma(x)\gamma'(x')$. It is obvious that the homomorphisms ψ and θ are mutually invertible.

The homomorphism $\gamma\otimes I$ is understood as follows. Let $f(a,\mathbf{e}) = \sum\limits_{k\geq 0}\sum\limits_{i_1,\ldots,i_k} a^{i_1\ldots i_k}\mathbf{e}_{i_1}\cdots\mathbf{e}_{i_k}$ be a general element of the algebra $\mathfrak{A}$, $a^{i_1\ldots i_k}\in\mathbb{F}$. If $f(a_1,\mathbf{e}_1;a_2,\mathbf{e}_2) = f_1(a_1,\mathbf{e}_1)f_2(a_2,\mathbf{e}_2)$ is an element of the algebra $\mathfrak{A}\hat{\otimes}\mathfrak{A}$, then

$$(\gamma\otimes If)(a_1,\mathbf{e}_1;a_2,\mathbf{e}_2;a_3,\mathbf{e}_3) = (\gamma f_1)(a_1,\mathbf{e}_1;a_2,\mathbf{e}_2)f_2(a_3,\mathbf{e}_3)$$

defines an action of the homomorphism $\gamma\otimes I$. The homomorphisms $I\otimes\gamma$, $I\otimes\varepsilon$, $I\otimes\bigcirc$ are defined analogously.

The diagrams (12)–(13) present the axioms of associativity, right unit and right inverse element, respectively. The first two diagrams show that γ and ε define on $\mathfrak{A}$ the structure of a co-algebra.

The conditions (12)–(14) coincide with axioms for classical formal groups [12]. The main difference from the classical case consists in the permutation conditions $\mathbf{e}_i\mathbf{e}_j = (-1)^{\deg\mathbf{e}_i\deg\mathbf{e}_j}\mathbf{e}_j\mathbf{e}_i$ for the unuts of the graded algebra $\mathfrak{A}$. As in the case of classical theory, one can establish a left analog of the equalities (13) and (14) [13].

Let us consider a supergroup $G(\mathfrak{A},\gamma,\bigcirc)$ over the field $\mathbb{F}=\mathbb{R}$. In this case we have $\mathfrak{A} = \mathcal{C}\ell_{p,q}$. The real Clifford algebra $\mathcal{C}\ell_{p,q}$ is central simple if $p-q\not\equiv 1,5 \pmod 8$. The graded tensor product of the two graded central simple algebras is also graded central simple [17, Theorem 2]. It is known that for the Clifford algebra with odd dimensionality, the isomorphisms are as follows: $\mathcal{C}\ell^+_{p,q+1}\simeq\mathcal{C}\ell_{p,q}$ and $\mathcal{C}\ell^+_{p+1,q}\simeq\mathcal{C}\ell_{q,p}$ [23, 24]. Thus, $\mathcal{C}\ell^+_{p,q+1}$ and $\mathcal{C}\ell^+_{p+1,q}$ are central simple algebras. Further, in accordance with Chevalley Theorem for the graded tensor product there is an isomorphism $\mathcal{C}\ell_{p,q}\hat{\otimes}\mathcal{C}\ell_{p',q'}\simeq\mathcal{C}\ell_{p+p',q+q'}$. Two algebras $\mathcal{C}\ell_{p,q}$ and $\mathcal{C}\ell_{p',q'}$ are said to be of the same class if $p+q'\equiv p'+q \pmod 8$. The graded central simple Clifford algebras over the field $\mathbb{F}=\mathbb{R}$ form eight similarity classes, which, as it is easy to see, coincide with the eight types of the algebras $\mathcal{C}\ell_{p,q}$. The set of these 8 types (classes) forms a Brauer-Wall group $BW_{\mathbb{R}}$ [17, 18]. It is obvious that an action of the supergroup $BW_{\mathbb{R}} = G(\mathcal{C}\ell_{p,q},\gamma,\bigcirc)$ has a cyclic structure, which is formally equivalent to the action of cyclic group $\mathbb{Z}_8$. The cyclic structure of the supergroup $BW_{\mathbb{R}}$ may be represented on the Trautman diagram (spinorial clock) [15, 16] (Fig. 1) by means of a transition $\mathcal{C}\ell^+_{p,q}\xrightarrow{h}\mathcal{C}\ell_{p,q}$ (the round on the diagram is realized by an hour-hand). At this point, the type of the algebra is defined on the diagram by an equality $q-p = h+8r$, where $h\in\{0,\ldots,7\}$, $r\in\mathbb{Z}$.



Fig.1 The Trautman diagram for the Brauer-Wall group $BW_{\mathbb{R}}$.



Fig.2 The Trautman diagram for the Brauer-Wall group $BW_{\mathbb{C}}$.

It is obvious that a group structure over $\mathcal{C}\!\ell_{p,q}$, defined by $BW_{\mathbb{R}}$, is related with the Atiyah-Bott-Shapiro periodicity [14]. In accordance with [14], the Clifford algebra over the field $\mathbb{F}=\mathbb{R}$ is modulo 8 periodic: $\mathcal{C}\!\ell_{p+8,q} \simeq \mathcal{C}\!\ell_{p,q} \otimes \mathcal{C}\!\ell_{8,0}$ ($\mathcal{C}\!\ell_{p,q+8} \simeq \mathcal{C}\!\ell_{p,q} \otimes \mathcal{C}\!\ell_{0,8}$).

Further, over the field $\mathbb{F}=\mathbb{C}$, there exist two types of the complex Clifford algebras: $\mathbb{C}_n$ and $\mathbb{C}_{n+1} \simeq \mathbb{C}_n \oplus \mathbb{C}_n$. Therefore, an action of $BW_{\mathbb{C}}$, defined on the set of these two types, is formally equivalent to the action of cyclic group $\mathbb{Z}_2$. The cyclic structure of the supergroup $BW_{\mathbb{C}} = G(\mathbb{C}_n, \gamma, \bigcirc)$ may be represented on the following Trautman diagram (Fig. 2) by means of a transition $\mathbb{C}^+_n \xrightarrow{h} \mathbb{C}_n$ (the round on the diagram is realized by an hour-hand). At this point, the type of the algebra on the diagram is defined by an equality $n = h + 2r$, where $h \in \{0,1\}$, $r \in \mathbb{Z}$. It is obvious that a group structure over $\mathbb{C}_n$, defined by the group $BW_{\mathbb{C}}$, is related immediately with a modulo 2 periodicity of the complex Clifford algebras [14, 22]: $\mathbb{C}_{n+2} \simeq \mathbb{C}_n \otimes \mathbb{C}_2$.

3.2 Superalgebras

As in the case of classical theory of groups and algebras, the supergroup $G(\mathfrak{A}, \gamma, \bigcirc)$ corresponds to a superalgebra (or $\mathbb{Z}_2$-graded Lie algebra). $\mathbb{Z}_2$-graded linear space over the field $\mathbb{F}$, endowed with a bilinear operation $[X_1, X_2]$, is called a *superalgebra.* At this point, the operation $[X_1, X_2]$ satisfies the following conditions:

$$
\begin{gathered}
[X_1, X_2] = -(-1)^{\deg(X_1)\deg(X_2)}\,[X_2, X_1]\,, \\
(-1)^{\deg(X_1)\deg(X_3)}\,[X_1,[X_2,X_3]] + (-1)^{\deg(X_2)\deg(X_1)}\,[X_2,[X_3,X_1]] + \\
+(-1)^{\deg(X_3)\deg(X_2)}\,[X_3,[X_1,X_2]] = 0, \\
\deg\left([X_1, X_2]\right) = \deg(X_1) + \deg(X_2).
\end{gathered}
$$

4 Clifford algebras and representations of $\mathbf{Spin}_+(1,3)$

Let us consider the field

$$
\boldsymbol{\psi}(\boldsymbol{\alpha}) = \langle x, \mathfrak{g}\,|\boldsymbol{\psi}\rangle, \tag{17}
$$

where $x \in T_4$, $\mathfrak{g} \in \mathbf{Spin}_+(1,3)$. The spinor group $\mathbf{Spin}_+(1,3) \simeq \mathrm{SU}(2) \otimes \mathrm{SU}(2)$ is a universal covering of the proper orthochronous Lorentz group $\mathrm{SO}_0(1,3)$. The parameters $x \in T_4$ and $\mathfrak{g} \in \mathbf{Spin}_+(1,3)$ describe position and orientation of the extended object defined by the field (17) (the field on the Poincaré group).

Theorem 3. *Let* $\mathbf{Spin}_+(1,3)$ *be the universal covering of the proper orthochronous Lorentz group* $\mathrm{SO}_0(1,3)$*. Then over* $\mathbb{F}=\mathbb{C}$ *the field* $(l,0) \oplus (0,\dot{l})$ *is constructed in the framework of a complex* *finite-dimensional representation* $\mathfrak{C}^{l_0+l_1-1,0}\oplus\mathfrak{C}^{0,l_0-l_1+1}$ *of* $\mathbf{Spin}_+(1,3)$ *defined on the spinspace* $\mathbb{S}_{2^k}\otimes\mathbb{S}_{2^r}$ *with the algebra*

$$\underbrace{\mathbb{C}_2\otimes\mathbb{C}_2\otimes\cdots\otimes\mathbb{C}_2}_{k\ times}\bigoplus\underbrace{\overset{*}{\mathbb{C}}_2\otimes\overset{*}{\mathbb{C}}_2\otimes\cdots\otimes\overset{*}{\mathbb{C}}_2}_{r\ times},$$

where $(l_0,l_1)=\left(\frac{k}{2},\frac{k}{2}+1\right)$, $(-l_0,l_1)=\left(-\frac{r}{2},\frac{r}{2}+1\right)$. *In turn, the field* $(l',l'')\oplus(\dot{l}'',\dot{l}')$ *is constructed in the framework of a representation* $\mathfrak{C}^{l_0+l_1-1,l_0-l_1+1}\oplus\mathfrak{C}^{l_0-l_1+1,l_0+l_1-1}$ *of* $\mathbf{Spin}_+(1,3)$ *defined on the spinspace* $\mathbb{S}_{2^{k+r}}\oplus\mathbb{S}_{2^{k+r}}$ *with the algebra*

$$\underbrace{\mathbb{C}_2\otimes\mathbb{C}_2\otimes\cdots\otimes\mathbb{C}_2\bigotimes\overset{*}{\mathbb{C}}_2\otimes\overset{*}{\mathbb{C}}_2\otimes\cdots\otimes\overset{*}{\mathbb{C}}_2}_{k+r\ times}\bigoplus\underbrace{\overset{*}{\mathbb{C}}_2\otimes\overset{*}{\mathbb{C}}_2\otimes\cdots\otimes\overset{*}{\mathbb{C}}_2\bigotimes\mathbb{C}_2\otimes\mathbb{C}_2\otimes\cdots\otimes\mathbb{C}_2}_{r+k\ times},$$

where $(l_0,l_1)=\left(\frac{k-r}{2},\frac{k+r}{2}+1\right)$. *In the case of* $\mathbb{F}=\mathbb{R}$ *the field* $\boldsymbol{\psi}(\boldsymbol{\alpha})$ *of type* $(l_0,0)$ *is constructed within real representations* $\mathfrak{M}^+=\{\mathfrak{R}^{l_0}_0,\mathfrak{R}^{l_0}_2,\mathfrak{H}^{l_0}_4,\mathfrak{H}^{l_0}_6\}$ *of* $\mathbf{Spin}_+(1,3)$ *defined on the spinspace* $\mathbb{S}_{2^r}$ *with the algebra*

$$\mathcal{C}\ell_{p,q}\simeq\underbrace{\mathcal{C}\ell_{s_i,t_j}\otimes\mathcal{C}\ell_{s_i,t_j}\otimes\cdots\otimes\mathcal{C}\ell_{s_i,t_j}}_{r\ times},$$

where $s_i,t_j\in\{0,1,2\}$. *In turn, the field* $\boldsymbol{\psi}(\boldsymbol{\alpha})$ *of type* $(l_0,0)\cup(0,l_0)$ *is constructed within representations* $\mathfrak{M}^-=\{\mathfrak{C}^{l_0}_{3,7},\mathfrak{R}^{l_0}_{0,2}\cup\mathfrak{R}^{l_0}_{0,2},\mathfrak{H}^{l_0}_{4,6}\cup\mathfrak{H}^{l_0}_{4,6}\}$ *of* $\mathbf{Spin}_+(1,3)$ *defined on the double spinspace* $\mathbb{S}_{2^r}\cup\mathbb{S}_{2^r}$ *with the algebra*

$$\mathcal{C}\ell_{p,q}\simeq\underbrace{\mathcal{C}\ell_{s_i,t_j}\otimes\mathcal{C}\ell_{s_i,t_j}\otimes\cdots\otimes\mathcal{C}\ell_{s_i,t_j}}_{r\ times}\bigcup\underbrace{\mathcal{C}\ell_{s_i,t_j}\otimes\mathcal{C}\ell_{s_i,t_j}\otimes\cdots\otimes\mathcal{C}\ell_{s_i,t_j}}_{r\ times}.$$

Proof. 1) Complex representations.
Since the spacetime algebra $\mathcal{C}\ell_{1,3}$ is the simple algebra, then the map (9) gives an irreducible representation of $\mathcal{C}\ell_{1,3}$ in the spinspace $\mathbb{S}_2(\mathbb{H})$. In turn, representations of the group $\mathbf{Spin}_+(1,3)\in\mathcal{C}\ell^+_{1,3}\simeq\mathcal{C}\ell_{3,0}$ are defined in the spinspace $\mathbb{S}_2(\mathbb{C})$.

Let us consider now spintensor representations of the group $\mathfrak{G}_+\simeq\mathrm{SL}(2;\mathbb{C})$ which, as is known, form the base of all the finite-dimensional representations of the Lorentz group, and also we consider their relationship with the complex Clifford algebras. From each complex Clifford algebra $\mathbb{C}_n=\mathbb{C}\otimes\mathcal{C}\ell_{p,q}$ $(n=p+q)$ we obtain the spinspace $\mathbb{S}_{2^{n/2}}$ which is a complexification of the minimal left ideal of the algebra $\mathcal{C}\ell_{p,q}$: $\mathbb{S}_{2^{n/2}}=\mathbb{C}\otimes I_{p,q}=\mathbb{C}\otimes\mathcal{C}\ell_{p,q}f_{pq}$, where f_{pq} is the primitive idempotent of the algebra $\mathcal{C}\ell_{p,q}$. Further, a spinspace related with the Pauli algebra $\mathbb{C}_2$ has the form $\mathbb{S}_2=\mathbb{C}\otimes I_{2,0}=\mathbb{C}\otimes\mathcal{C}\ell_{2,0}f_{20}$ or $\mathbb{S}_2=\mathbb{C}\otimes I_{1,1}=\mathbb{C}\otimes\mathcal{C}\ell_{1,1}f_{11}(\mathbb{C}\otimes I_{0,2}=\mathbb{C}\otimes\mathcal{C}\ell_{0,2}f_{02})$. Therefore, the tensor product of the k algebras $\mathbb{C}_2$ induces a tensor product of the k spinspaces $\mathbb{S}_2$:

$$\mathbb{S}_2\otimes\mathbb{S}_2\otimes\cdots\otimes\mathbb{S}_2=\mathbb{S}_{2^k}.$$

Vectors of the spinspace $\mathbb{S}_{2^k}$ (or elements of the minimal left ideal of $\mathbb{C}_{2k}$) are spintensors of the following form:

$$\boldsymbol{s}^{\alpha_1\alpha_2\cdots\alpha_k}=\sum\boldsymbol{s}^{\alpha_1}\otimes\boldsymbol{s}^{\alpha_2}\otimes\cdots\otimes\boldsymbol{s}^{\alpha_k},\tag{18}$$

where summation is produced on all the index collections $(\alpha_1\ldots\alpha_k)$, $\alpha_i=1,2$. For the each spinor $\boldsymbol{s}^{\alpha_i}$ from (18) we have ${}'\boldsymbol{s}^{\alpha'_i}=\sigma^{\alpha'_i}_{\alpha_i}\boldsymbol{s}^{\alpha_i}$. Therefore, in general case we obtain

$${}'\boldsymbol{s}^{\alpha'_1\alpha'_2\cdots\alpha'_k}=\sum\sigma^{\alpha'_1}_{\alpha_1}\sigma^{\alpha'_2}_{\alpha_2}\cdots\sigma^{\alpha'_k}_{\alpha_k}\boldsymbol{s}^{\alpha_1\alpha_2\cdots\alpha_k}.\tag{19}$$

A representation (19) is called *undotted spintensor representation of the proper Lorentz group of the rank k.*

Further, let $\overset{*}{\mathbb{C}}_2$ be the Pauli algebra with the coefficients which are complex conjugate to the coefficients of $\mathbb{C}_2$. Let us show that the algebra $\overset{*}{\mathbb{C}}_2$ is derived from $\mathbb{C}_2$ under action of the automorphism $\mathcal{A} \rightarrow \mathcal{A}^\star$ or antiautomorphism $\mathcal{A} \rightarrow \widetilde{\mathcal{A}}$ (see the definitions (5) and (7)). Indeed, in virtue of an isomorphism $\mathbb{C}_2 \simeq \mathcal{C}\!\ell_{3,0}$ a general element

$$\mathcal{A} = a^0\mathbf{e}_0 + \sum_{i=1}^{3} a^i\mathbf{e}_i + \sum_{i=1}^{3}\sum_{j=1}^{3} a^{ij}\mathbf{e}_{ij} + a^{123}\mathbf{e}_{123}$$

of the algebra $\mathcal{C}\!\ell_{3,0}$ can be written in the form

$$\mathcal{A} = (a^0 + \omega a^{123})\mathbf{e}_0 + (a^1 + \omega a^{23})\mathbf{e}_1 + (a^2 + \omega a^{31})\mathbf{e}_2 + (a^3 + \omega a^{12})\mathbf{e}_3, \tag{20}$$

where $\omega = \mathbf{e}_{123}$. Since ω belongs to a center of the algebra $\mathcal{C}\!\ell_{3,0}$ (ω commutes with all the basis elements) and $\omega^2 = -1$, then we can to suppose $\omega \equiv i$. The action of the automorphism $\star$ on the homogeneous element $\mathcal{A}$ of the degree k is defined by the formula (5). In accordance with this the action of the automorphism $\mathcal{A} \rightarrow \mathcal{A}^\star$, where $\mathcal{A}$ is the element (20), has the form

$$\mathcal{A} \longrightarrow \mathcal{A}^\star = -(a^0 - \omega a^{123})\mathbf{e}_0 - (a^1 - \omega a^{23})\mathbf{e}_1 - (a^2 - \omega a^{31})\mathbf{e}_2 - (a^3 - \omega a^{12})\mathbf{e}_3. \tag{21}$$

Therefore, $\star : \mathbb{C}_2 \rightarrow -\overset{*}{\mathbb{C}}_2$. Correspondingly, the action of the antiautomorphism $\mathcal{A} \rightarrow \widetilde{\mathcal{A}}$ on the homogeneous element $\mathcal{A}$ of the degree k is defined by the formula (7). Thus, for the element (20) we obtain

$$\mathcal{A} \longrightarrow \widetilde{\mathcal{A}} = (a^0 - \omega a^{123})\mathbf{e}_0 + (a^1 - \omega a^{23})\mathbf{e}_1 + (a^2 - \omega a^{31})\mathbf{e}_2 + (a^3 - \omega a^{12})\mathbf{e}_3, \tag{22}$$

that is, $\widetilde{\ } : \mathbb{C}_2 \rightarrow \overset{*}{\mathbb{C}}_2$. This allows us to define an algebraic analogue of the Wigner's representation doubling: $\mathbb{C}_2 \oplus \overset{*}{\mathbb{C}}_2$. Further, from (20) it follows that $\mathcal{A} = \mathcal{A}_1 + \omega\mathcal{A}_2 = (a^0\mathbf{e}_0 + a^1\mathbf{e}_1 + a^2\mathbf{e}_2 + a^3\mathbf{e}_3) + \omega(a^{123}\mathbf{e}_0 + a^{23}\mathbf{e}_1 + a^{31}\mathbf{e}_2 + a^{12}\mathbf{e}_3)$. In general case, by virtue of an isomorphism $\mathbb{C}_{2k} \simeq \mathcal{C}\!\ell_{p,q}$, where $\mathcal{C}\!\ell_{p,q}$ is a real Clifford algebra with a division ring $\mathbb{K} \simeq \mathbb{C}$, $p - q \equiv 3, 7 \pmod 8$, we have for the general element of $\mathcal{C}\!\ell_{p,q}$ an expression $\mathcal{A} = \mathcal{A}_1 + \omega\mathcal{A}_2$, here $\omega^2 = \mathbf{e}^2_{12\ldots p+q} = -1$ and, therefore, $\omega \equiv i$. Thus, from $\mathbb{C}_{2k}$ under action of the automorphism $\mathcal{A} \rightarrow \mathcal{A}^\star$ we obtain a general algebraic doubling

$$\mathbb{C}_{2k} \oplus \overset{*}{\mathbb{C}}_{2k}. \tag{23}$$

The tensor product $\overset{*}{\mathbb{C}}_2 \otimes \overset{*}{\mathbb{C}}_2 \otimes \cdots \otimes \overset{*}{\mathbb{C}}_2 \simeq \overset{*}{\mathbb{C}}_{2r}$ of the r algebras $\overset{*}{\mathbb{C}}_2$ induces the tensor product of the r spinspaces $\dot{\mathbb{S}}_2$:

$$\dot{\mathbb{S}}_2 \otimes \dot{\mathbb{S}}_2 \otimes \cdots \otimes \dot{\mathbb{S}}_2 = \dot{\mathbb{S}}_{2^r}.$$

Vectors of the spinspace $\dot{\mathbb{S}}_{2^r}$ have the form

$$\boldsymbol{s}^{\dot{\alpha}_1\dot{\alpha}_2\cdots\dot{\alpha}_r} = \sum \boldsymbol{s}^{\dot{\alpha}_1} \otimes \boldsymbol{s}^{\dot{\alpha}_2} \otimes \cdots \otimes \boldsymbol{s}^{\dot{\alpha}_r}, \tag{24}$$

where the each cospinor $\boldsymbol{s}^{\dot{\alpha}_i}$ from (24) is transformed by the rule ${}'\boldsymbol{s}^{\dot{\alpha}'_i} = \sigma^{\dot{\alpha}'_i}_{\dot{\alpha}_i}\boldsymbol{s}^{\dot{\alpha}_i}$. Therefore,

$${}'\boldsymbol{s}^{\dot{\alpha}'_1\dot{\alpha}'_2\cdots\dot{\alpha}'_r} = \sum \sigma^{\dot{\alpha}'_1}_{\dot{\alpha}_1}\sigma^{\dot{\alpha}'_2}_{\dot{\alpha}_2}\cdots\sigma^{\dot{\alpha}'_r}_{\dot{\alpha}_r}\boldsymbol{s}^{\dot{\alpha}_1\dot{\alpha}_2\cdots\dot{\alpha}_r}. \tag{25}$$

The representation (25) is called *a dotted spintensor representation of the proper Lorentz group of the rank r.*

In general case we have a tensor product of the k algebras $\mathbb{C}_2$ and the r algebras $\overset{*}{\mathbb{C}}_2$:

$$\mathbb{C}_2\otimes\mathbb{C}_2\otimes\cdots\otimes\mathbb{C}_2\bigotimes\overset{*}{\mathbb{C}}_2\otimes\overset{*}{\mathbb{C}}_2\otimes\cdots\otimes\overset{*}{\mathbb{C}}_2\simeq\mathbb{C}_{2k}\otimes\overset{*}{\mathbb{C}}_{2r},$$

which induces a spinspace

$$\mathbb{S}_2\otimes\mathbb{S}_2\otimes\cdots\otimes\mathbb{S}_2\bigotimes\dot{\mathbb{S}}_2\otimes\dot{\mathbb{S}}_2\otimes\cdots\otimes\dot{\mathbb{S}}_2=\mathbb{S}_{2^{k+r}}$$

with the vectors

$$\boldsymbol{s}^{\alpha_1\alpha_2\cdots\alpha_k\dot{\alpha}_1\dot{\alpha}_2\cdots\dot{\alpha}_r}=\sum\boldsymbol{s}^{\alpha_1}\otimes\boldsymbol{s}^{\alpha_2}\otimes\cdots\otimes\boldsymbol{s}^{\alpha_k}\otimes\boldsymbol{s}^{\dot{\alpha}_1}\otimes\boldsymbol{s}^{\dot{\alpha}_2}\otimes\cdots\otimes\boldsymbol{s}^{\dot{\alpha}_r}. \tag{26}$$

In this case we have a natural unification of the representations (19) and (25):

$${}'\boldsymbol{s}^{\alpha'_1\alpha'_2\cdots\alpha'_k\dot{\alpha}'_1\dot{\alpha}'_2\cdots\dot{\alpha}'_r}=\sum\sigma^{\alpha'_1}_{\alpha_1}\sigma^{\alpha'_2}_{\alpha_2}\cdots\sigma^{\alpha'_k}_{\alpha_k}\sigma^{\dot{\alpha}'_1}_{\dot{\alpha}_1}\sigma^{\dot{\alpha}'_2}_{\dot{\alpha}_2}\cdots\sigma^{\dot{\alpha}'_r}_{\dot{\alpha}_r}\boldsymbol{s}^{\alpha_1\alpha_2\cdots\alpha_k\dot{\alpha}_1\dot{\alpha}_2\cdots\dot{\alpha}_r}. \tag{27}$$

So, a representation (27) is called *a spintensor representation of the proper Lorentz group of the rank* (k,r).

Further, let $\mathfrak{g}\rightarrow T_{\mathfrak{g}}$ be an arbitrary linear representation of the proper orthochronous Lorentz group $\mathfrak{G}_+=\mathrm{SO}_0(1,3)$ and let $\mathsf{A}_i(t)=T_{a_i(t)}$ be an infinitesimal operator corresponding to the rotation $a_i(t)\in\mathfrak{G}_+$. Analogously, let $\mathsf{B}_i(t)=T_{b_i(t)}$, where $b_i(t)\in\mathfrak{G}_+$ is the hyperbolic rotation. The operators A_i and B_i satisfy to the following relations:

$$\left.\begin{array}{lll}
[\mathsf{A}_1,\mathsf{A}_2]=\mathsf{A}_3, & [\mathsf{A}_2,\mathsf{A}_3]=\mathsf{A}_1, & [\mathsf{A}_3,\mathsf{A}_1]=\mathsf{A}_2,\\
[\mathsf{B}_1,\mathsf{B}_2]=-\mathsf{A}_3, & [\mathsf{B}_2,\mathsf{B}_3]=-\mathsf{A}_1, & [\mathsf{B}_3,\mathsf{B}_1]=-\mathsf{A}_2,\\
[\mathsf{A}_1,\mathsf{B}_1]=0, & [\mathsf{A}_2,\mathsf{B}_2]=0, & [\mathsf{A}_3,\mathsf{B}_3]=0,\\
[\mathsf{A}_1,\mathsf{B}_2]=\mathsf{B}_3, & [\mathsf{A}_1,\mathsf{B}_3]=-\mathsf{B}_2, & \\
[\mathsf{A}_2,\mathsf{B}_3]=\mathsf{B}_1, & [\mathsf{A}_2,\mathsf{B}_1]=-\mathsf{B}_3, & \\
[\mathsf{A}_3,\mathsf{B}_1]=\mathsf{B}_2, & [\mathsf{A}_3,\mathsf{B}_2]=-\mathsf{B}_1. &
\end{array}\right\} \tag{28}$$

As is known [25], finite-dimensional (spinor) representations of the group $\mathrm{SO}_0(1,3)$ in the space of symmetrical polynomials $\mathrm{Sym}_{(k,r)}$ have the following form:

$$T_{\mathfrak{g}}q(\xi,\overline{\xi})=(\gamma\xi+\delta)^{l_0+l_1-1}\overline{(\gamma\xi+\delta)}^{l_0-l_1+1}q\left(\frac{\alpha\xi+\beta}{\gamma\xi+\delta};\frac{\overline{\alpha\xi+\beta}}{\overline{\gamma\xi+\delta}}\right), \tag{29}$$

where $k=l_0+l_1-1$, $r=l_0-l_1+1$, and the pair (l_0,l_1) defines some representation of the group $\mathrm{SO}_0(1,3)$ in the Gel'fand-Naimark basis:

$$H_3\xi_{k\nu}=m\xi_{k\nu},$$

$$H_+\xi_{k\nu}=\sqrt{(k+\nu+1)(k-\nu)}\,\xi_{k,\nu+1},$$

$$H_-\xi_{k\nu}=\sqrt{(k+\nu)(k-\nu+1)}\,\xi_{k,\nu-1},$$

$$F_3\xi_{k\nu}=C_l\sqrt{k^2-\nu^2}\,\xi_{k-1,\nu}-A_l\nu\xi_{k,\nu}-C_{k+1}\sqrt{(k+1)^2-\nu^2}\,\xi_{k+1,\nu},$$

$$F_+\xi_{k\nu}=C_k\sqrt{(k-\nu)(k-\nu-1)}\,\xi_{k-1,\nu+1}-A_k\sqrt{(k-\nu)(k+\nu+1)}\,\xi_{k,\nu+1}+$$
$$+C_{k+1}\sqrt{(k+\nu+1)(k+\nu+2)}\,\xi_{k+1,\nu+1},$$

$$F_-\xi_{k\nu} = -C_k\sqrt{(k+\nu)(k+\nu-1)}\xi_{k-1,\nu-1} - A_k\sqrt{(k+\nu)(k-\nu+1)}\xi_{k,\nu-1} - \\ - C_{k+1}\sqrt{(k-\nu+1)(k-\nu+2)}\xi_{k+1,\nu-1},$$

$$A_k = \frac{il_0l_1}{k(k+1)}, \quad C_k = \frac{i}{k}\sqrt{\frac{(k^2-l_0^2)(k^2-l_1^2)}{4k^2-1}}, \tag{30}$$
$$\nu = -k, -k+1, \dots, k-1, k,$$
$$k = l_0\,, l_0+1, \dots,$$

where l_0 is positive integer or half-integer number, l_1 is an arbitrary complex number. These formulae define a finite-dimensional representation of the group $\mathrm{SO}_0(1,3)$ when $l_1^2 = (l_0+p)^2$, p is some natural number. In the case $l_1^2 \neq (l_0+p)^2$ we have an infinite-dimensional representation of $\mathrm{SO}_0(1,3)$. The operators $H_3, H_+, H_-, F_3, F_+, F_-$ are

$$H_+ = i\mathsf{A}_1 - \mathsf{A}_2, \quad H_- = i\mathsf{A}_1 + \mathsf{A}_2, \quad H_3 = i\mathsf{A}_3,$$
$$F_+ = i\mathsf{B}_1 - \mathsf{B}_2, \quad F_- = i\mathsf{B}_1 + \mathsf{B}_2, \quad F_3 = i\mathsf{B}_3.$$

Let us consider the operators

$$\mathsf{X}_l = \frac{1}{2}i(\mathsf{A}_l + i\mathsf{B}_l), \quad \mathsf{Y}_l = \frac{1}{2}i(\mathsf{A}_l - \mathbf{i}\mathsf{B}_l), \tag{31}$$
$$(l = 1,2,3).$$

Using the relations (28), we find that

$$[\mathsf{X}_k, \mathsf{X}_l] = i\varepsilon_{klm}\mathsf{X}_m, \quad [\mathsf{Y}_l, \mathsf{Y}_m] = i\varepsilon_{lmn}\mathsf{Y}_n, \quad [\mathsf{X}_l, \mathsf{Y}_m] = 0. \tag{32}$$

Further, introducing generators of the form

$$\left.\begin{array}{l} \mathsf{X}_+ = \mathsf{X}_1 + i\mathsf{X}_2, \quad \mathsf{X}_- = \mathsf{X}_1 - i\mathsf{X}_2, \\ \mathsf{Y}_+ = \mathsf{Y}_1 + i\mathsf{Y}_2, \quad \mathsf{Y}_- = \mathsf{Y}_1 - i\mathsf{Y}_2, \end{array}\right\} \tag{33}$$

we see that in virtue of commutativity of the relations (32) a space of an irreducible finite–dimensional representation of the group $\mathrm{SL}(2,\mathbb{C})$ can be spanned on the totality of $(2l+1)(2\dot{l}+1)$ basis vectors $\mid l,m;\dot{l},\dot{m}\rangle$, where $l,m,\dot{l},\dot{m}$ are integer or half–integer numbers, $-l \leq m \leq l$, $-\dot{l} \leq \dot{m} \leq \dot{l}$. Therefore,

$$\begin{aligned}
&\mathsf{X}_- \mid l,m;\dot{l},\dot{m}\rangle = \sqrt{(l+m)(l-m+1)} \mid l,m-1,\dot{l},\dot{m}\rangle \;\; (m > -l),\\
&\mathsf{X}_+ \mid l,m;\dot{l},\dot{m}\rangle = \sqrt{(l-m)(l+m+1)} \mid l,m+1;\dot{l},\dot{m}\rangle \;\; (m < l),\\
&\mathsf{X}_3 \mid l,m;\dot{l},\dot{m}\rangle = m \mid l,m;\dot{l},\dot{m}\rangle,\\
&\mathsf{Y}_- \mid l,m;\dot{l},\dot{m}\rangle = \sqrt{(\dot{l}+\dot{m})(\dot{l}-\dot{m}+1)} \mid l,m;\dot{l},\dot{m}-1\rangle \;\; (\dot{m} > -\dot{l}),\\
&\mathsf{Y}_+ \mid l,m;\dot{l},\dot{m}\rangle = \sqrt{(\dot{l}-\dot{m})(\dot{l}+\dot{m}+1)} \mid l,m;\dot{l},\dot{m}+1\rangle \;\; (\dot{m} < \dot{l}),\\
&\mathsf{Y}_3 \mid l,m;\dot{l},\dot{m}\rangle = \dot{m} \mid l,m;\dot{l},\dot{m}\rangle.
\end{aligned} \tag{34}$$

From the relations (32) it follows that each of the sets of infinitesimal operators X and Y generates the group $\mathrm{SU}(2)$ and these two groups commute with each other. Thus, from the relations (32) and (34) it follows that the group $\mathrm{SL}(2,\mathbb{C})$, in essence, is equivalent locally to the group $\mathrm{SU}(2)\otimes\mathrm{SU}(2)$. This representation for the Lorentz group was first given by Van der Waerden in [26]. The following relations between generators $\mathsf{Y}_{\pm}$, $\mathsf{X}_{\pm}$, Y_3, X_3 and $H_{\pm}$, $F_{\pm}$, H_3, F_3 define a relationship between the Van der Waerden and Gel'fand-Naimark bases:

$$\begin{array}{rclrcl}
\mathsf{Y}_{+} & = & -\frac{1}{2}(F_{+}+iH_{+}), & \mathsf{X}_{+} & = & \frac{1}{2}(F_{+}-iH_{+}), \\
\mathsf{Y}_{-} & = & -\frac{1}{2}(F_{-}+iH_{-}), & \mathsf{X}_{-} & = & \frac{1}{2}(F_{-}-iH_{-}), \\
\mathsf{Y}_{3} & = & -\frac{1}{2}(F_{3}+iH_{3}), & \mathsf{X}_{3} & = & \frac{1}{2}(F_{3}-iH_{3}).
\end{array}$$

The relation between the numbers l_0, l_1 and the number l (the weight of representation in the basis (34)) is given by the following formula:

$$(l_0, l_1) = (l, l+1)\,.$$

Whence it immediately follows that

$$l = \frac{l_0 + l_1 - 1}{2}. \tag{35}$$

As is known [25], if an irreducible representation of the proper Lorentz group $\mathrm{SO}_0(1,3)$ is defined by the pair (l_0, l_1), then a conjugated representation is also irreducible and is defined by a pair $\pm(l_0, -l_1)$. Therefore,

$$(l_0, l_1) = \left(-\dot{l},\, \dot{l}+1\right).$$

Thus,

$$\dot{l} = \frac{l_0 - l_1 + 1}{2}. \tag{36}$$

Further, representations $\mathfrak{C}^{s_1,s_2}$ and $\mathfrak{C}^{s'_1,s'_2}$ are called *interlocking irreducible representations of the Lorentz group* , that is, such representations that $s'_1 = s_1 \pm 1$, $s'_2 = s_2 \pm 1$ [27]. The two most full schemes of the interlocking irreducible representations of the Lorentz group (Gel'fand-Yaglom chains) for integer and half-integer spins are shown on the Fig. 3 and Fig. 4. As follows from Fig. 3 the simplest field is the scalar field $\mathfrak{C}^{0,0}$. This field is described by the Fock-Klein-Gordon equation. In its turn, the simplest field from the Fermi-scheme (Fig. 4) is the electron-positron (spinor) field corresponding to the following interlocking scheme:

$$\mathfrak{C}^{1,0} \longleftrightarrow \mathfrak{C}^{0,-1}\;.$$

This field is described by the Dirac equation. Further, the next field from the Bose-scheme (Fig. 3) is a photon field (Maxwell field) defined within the interlocking scheme

$$\mathfrak{C}^{2,0} \longleftrightarrow \mathfrak{C}^{1,-1} \longleftrightarrow \mathfrak{C}^{0,-2}\;.$$

This interlocking scheme leads to the Maxwell equations. The fields $(1/2,0)\oplus(0,1/2)$ and $(1,0)\oplus(0,1)$ (Dirac and Maxwell fields) are particular cases of fields of the type $(l,0)\oplus(0,l)$. Wave equations for such fields and their general solutions were found in the works [28, 29, 30].

It is easy to see that the interlocking scheme, corresponded to the Maxwell field, contains the field of tensor type, $\mathfrak{C}^{1,-1}$. Further, the next interlocking scheme (see Fig. 4)

$$\mathfrak{C}^{3,0} \longleftrightarrow \mathfrak{C}^{2,-1} \longleftrightarrow \mathfrak{C}^{1,-2} \longleftrightarrow \mathfrak{C}^{0,-3}\;,$$

corresponding to the Pauli-Fierz equations [31], contains a chain of the type

$$\mathfrak{C}^{2,-1} \longleftrightarrow \mathfrak{C}^{1,-2}\;.$$

$$\begin{tikzcd}
 & & & & \mathfrak{C}^{s,0} \arrow[r, dash] \arrow[d, dash] & \cdots \\
 & & & & \vdots \arrow[d, dash] & \\
 & & \mathfrak{C}^{4,0} \arrow[r, dash] \arrow[d, dash] & \cdots \arrow[r, dash] & \mathfrak{C}^{s+2,2-s} \arrow[r, dash] \arrow[d, dash] & \cdots \\
 & \mathfrak{C}^{2,0} \arrow[r, dash] \arrow[d, dash] & \mathfrak{C}^{3,-1} \arrow[r, dash] \arrow[d, dash] & \cdots \arrow[r, dash] & \mathfrak{C}^{s+1,1-s} \arrow[r, dash] \arrow[d, dash] & \cdots \\
\mathfrak{C}^{0,0} \arrow[r, dash] & \mathfrak{C}^{1,-1} \arrow[r, dash] \arrow[d, dash] & \mathfrak{C}^{2,-2} \arrow[r, dash] \arrow[d, dash] & \cdots \arrow[r, dash] & \mathfrak{C}^{s,-s} \arrow[r, dash] \arrow[d, dash] & \cdots \\
 & \mathfrak{C}^{0,-2} \arrow[r, dash] & \mathfrak{C}^{1,-3} \arrow[r, dash] \arrow[d, dash] & \cdots \arrow[r, dash] & \mathfrak{C}^{s-1,-1-s} \arrow[r, dash] \arrow[d, dash] & \cdots \\
 & & \mathfrak{C}^{0,-4} \arrow[r, dash] & \cdots \arrow[r, dash] & \mathfrak{C}^{s-2,-2-s} \arrow[r, dash] \arrow[d, dash] & \cdots \\
 & & & & \vdots \arrow[d, dash] & \\
 & & & & \mathfrak{C}^{0,-2s} \arrow[r, dash] & \cdots
\end{tikzcd}$$

Fig. 3: Complex interlocking representations of $\mathbf{Spin}_+(1,3)$ for the fields of integer spin (Bose-scheme).

In such a way we come to wave equations for the fields $\boldsymbol{\psi}(\boldsymbol{\alpha}) = \langle x, \mathfrak{g}\,|\boldsymbol{\psi}\rangle$ of tensor type $(l_1, l_2) \oplus (l_2, l_1)$. Wave equations for such fields and their general solutions were found in the work [32].

A relation between the numbers l_0, l_1 of the Gel'fand-Naimark representation (30) and the number k of the factors $\mathbb{C}_2$ in the product $\mathbb{C}_2 \otimes \mathbb{C}_2 \otimes \cdots \otimes \mathbb{C}_2$ is given by the following formula:

$$(l_0, l_1) = \left(\frac{k}{2}, \frac{k}{2} + 1\right),$$

Hence it immediately follows that $k = l_0 + l_1 - 1$. Thus, we have *a complex representation* $\mathfrak{C}^{l_0+l_1-1,0}$ *of the group* $\mathbf{Spin}_+(1,3)$ *in the spinspace* $\mathbb{S}_{2^k}$. If the representation $\mathfrak{C}^{l_0+l_1-1,0}$ is reducible, then the space $\mathbb{S}_{2^k}$ is decomposed into a direct sum of irreducible subspaces, that is, it is possible to choose in $\mathbb{S}_{2^k}$ such a basis, in which all the matrices take a block-diagonal form. Then the field $\boldsymbol{\psi}(\boldsymbol{\alpha})$ is reduced to some number of the fields corresponding to irreducible representations of the group $\mathbf{Spin}_+(1,3)$, each of which is transformed independently from the other, and the field $\boldsymbol{\psi}(\boldsymbol{\alpha})$ in this case is a collection of the fields with more simple structure. It is obvious that these more simple fields correspond to irreducible representations $\mathfrak{C}$.

Analogously, a relation between the numbers l_0, l_1 of the Gel'fand-Naimark representation (30) and the number r of the factors $\overset{*}{\mathbb{C}}_2$ in the product $\overset{*}{\mathbb{C}}_2 \otimes \overset{*}{\mathbb{C}}_2 \otimes \cdots \otimes \overset{*}{\mathbb{C}}_2$ is given by the following formula:

$$(-l_0, l_1) = \left(-\frac{r}{2}, \frac{r}{2} + 1\right).$$

Hence it immediately follows that $r = l_0 - l_1 + 1$. Thus, we have a complex representation $\mathfrak{C}^{0,l_0-l_1+1}$ of $\mathbf{Spin}_+(1,3)$ in the spinspace $\mathbb{S}_{2^r}$.

As is known [33, 25, 34], a system of irreducible finite-dimensional representations of the group $\mathfrak{G}_+$ is realized in the space $\mathrm{Sym}_{(k,r)} \subset \mathbb{S}_{2^{k+r}}$ of symmetric spintensors. The dimensionality of

$$\begin{tikzcd}
 & & & \mathfrak{C}^{2s,0} \arrow[r, dash] \arrow[d, dash] & \cdots \\
 & & & \vdots \arrow[d, dash] & \\
 & \mathfrak{C}^{3,0} \arrow[r, dash] \arrow[d, dash] & \cdots \arrow[r, dash] & \mathfrak{C}^{(2s+3)/2,(3-2s)2} \arrow[r, dash] \arrow[d, dash] & \cdots \\
\mathfrak{C}^{1,0} \arrow[r, dash] \arrow[d, dash] & \mathfrak{C}^{2,-1} \arrow[r, dash] \arrow[d, dash] & \cdots \arrow[r, dash] & \mathfrak{C}^{(2s+1)/2,(1-2s)/2} \arrow[r, dash] \arrow[d, dash] & \cdots \\
\mathfrak{C}^{0,-1} \arrow[r, dash] & \mathfrak{C}^{1,-2} \arrow[r, dash] \arrow[d, dash] & \cdots \arrow[r, dash] & \mathfrak{C}^{(2s-1)/2,-(2s+1)/2} \arrow[r, dash] \arrow[d, dash] & \cdots \\
 & \mathfrak{C}^{0,-3} \arrow[r, dash] & \cdots \arrow[r, dash] & \mathfrak{C}^{(2s-3)/2,-(2s+3)/2} \arrow[r, dash] \arrow[d, dash] & \cdots \\
 & & & \vdots \arrow[d, dash] & \\
 & & & \mathfrak{C}^{0,-2s} \arrow[r, dash] & \cdots
\end{tikzcd}$$

Fig. 4: Complex interlocking representations of $\mathbf{Spin}_+(1,3)$ for the fields of half-integer spin (Fermi-scheme).

$\mathrm{Sym}_{(k,r)}$ is equal to $(k+1)(r+1)$. A representation of the group $\mathfrak{G}_+$, defined by such spintensors, is irreducible and denoted by the symbol $\mathfrak{D}^{(l,\dot{l})}(\sigma)$, where $2l=k$, $2\dot{l}=r$, the numbers l and $\dot{l}$ are integer or half-integer. In general case, the field $\boldsymbol{\psi}(\boldsymbol{\alpha})$ is the field of type $(l,\dot{l})$. As a rule, in physics there are three basic types of the fields.

a) The field of type $(l,0)$. The structure of this field (or the field $(0,\dot{l})$) is described by the representation $\mathfrak{D}^{(l,0)}(\sigma)$ ($\mathfrak{D}^{(0,\dot{l})}(\sigma)$), which is realized in the space $\mathbb{S}_{2^k}$ ($\mathbb{S}_{2^r}$). At this point, the algebra $\mathbb{C}_{2k}\simeq\mathbb{C}_2\otimes\mathbb{C}_2\otimes\cdots\otimes\mathbb{C}_2$ (correspondingly, $\overset{*}{\mathbb{C}}_{2k}\simeq\overset{*}{\mathbb{C}}_2\otimes\overset{*}{\mathbb{C}}_2\otimes\cdots\otimes\overset{*}{\mathbb{C}}_2$) is associated with the field of the type $(l,0)$ (correspondingly, $(0,\dot{l})$). The trivial case $l=0$ corresponds to *a Pauli-Weisskopf field* describing the scalar particles. Further, at $l=\dot{l}=1/2$ we have *a Weyl field* describing the neutrino. At this point, the antineutrino is described by a fundamental representation $\mathfrak{D}^{(1/2,0)}(\sigma)=\sigma$ of the group $\mathfrak{G}_+$ and the algebra $\mathbb{C}_2$. Correspondingly, the neutrino is described by a conjugated representation $\mathfrak{D}^{(0,1/2)}(\sigma)$ and the algebra $\overset{*}{\mathbb{C}}_2$. In essence, one can say that the algebra $\mathbb{C}_2$ ($\overset{*}{\mathbb{C}}_2$) is the basic building block, from which other physical fields built by means of direct sum or tensor product. One can say also that this situation looks like the de Broglie fusion method [35].

b) The field of type $(l,0)\oplus(0,\dot{l})$. The structure of this field admits a space inversion and, therefore, in accordance with a Wigner's doubling [36] is described by a representation $\mathfrak{D}^{(l,0)}\oplus\mathfrak{D}^{(0,\dot{l})}$ of the group $\mathfrak{G}_+$. This representation is realized in the space $\mathbb{S}_{2^{2k}}$. The Clifford algebra, related with this representation, is a direct sum $\mathbb{C}_{2k}\oplus\overset{*}{\mathbb{C}}_{2k}\simeq\mathbb{C}_2\otimes\mathbb{C}_2\otimes\cdots\otimes\mathbb{C}_2\bigoplus\overset{*}{\mathbb{C}}_2\otimes\overset{*}{\mathbb{C}}_2\otimes\cdots\otimes\overset{*}{\mathbb{C}}_2$. In the simplest case $l=1/2$ we have *bispinor (electron–positron) Dirac field* $(1/2,0)\oplus(0,1/2)$ with the algebra $\mathbb{C}_2\oplus\overset{*}{\mathbb{C}}_2$. It should be noted that the Dirac algebra $\mathbb{C}_4$, considered as a tensor product

$\mathbb{C}_2 \otimes \mathbb{C}_2$ (or $\mathbb{C}_2 \otimes \overset{*}{\mathbb{C}}_2$) in accordance with (18) (or (26)) gives rise to spintensors $\boldsymbol{s}^{\alpha_1\alpha_2}$ (or $\boldsymbol{s}^{\alpha_1\dot{\alpha}_1}$), but it contradicts with the usual definition of the Dirac bispinor as a pair $(\boldsymbol{s}^{\alpha_1}, \boldsymbol{s}^{\dot{\alpha}_1})$. Therefore, the Clifford algebra, associated with the Dirac field, is $\mathbb{C}_2 \oplus \overset{*}{\mathbb{C}}_2$, and a spinspace of this sum in virtue of unique decomposition $\mathbb{S}_2 \oplus \dot{\mathbb{S}}_2 = \mathbb{S}_4$ is a spinspace of $\mathbb{C}_4$.

c) Tensor fields $(l', l'') \oplus (\dot{l}'', \dot{l}')$. The fields (l', l'') and $(\dot{l}'', \dot{l}')$ are defined within the arbitrary spin chains (see Fig. 3 and Fig. 4). Universal coverings of these spin chains are constructed within the representations $\mathfrak{C}^{l_0+l_1-1,l_0-l_1+1}$ and $\mathfrak{C}^{l_0-l_1+1,l_0+l_1-1}$ of $\mathbf{Spin}_+(1,3)$ in the spinspaces $\mathbb{S}_{2^{k+r}}$ associated with the algebra $\mathbb{C}_2 \otimes \mathbb{C}_2 \otimes \cdots \otimes \mathbb{C}_2 \bigotimes \overset{*}{\mathbb{C}}_2 \otimes \overset{*}{\mathbb{C}}_2 \otimes \cdots \otimes \overset{*}{\mathbb{C}}_2$. A relation between the numbers l_0, l_1 of the Gel'fand-Naimark basis (30) and the numbers k and r of the factors $\mathbb{C}_2$ and $\overset{*}{\mathbb{C}}_2$ is given by the following formula:

$$(l_0, l_1) = \left(\frac{k-r}{2}, \frac{k+r}{2} + 1\right).$$

2) Real representations.

As is known [25], if an irreducible representation of the proper Lorentz group $\mathrm{SO}_0(1,3)$ is defined by the pair (l_0, l_1), then a conjugated representation is also irreducible and is defined by a pair $\pm(l_0, -l_1)$. Hence it follows that an irreducible representation is equivalent to its conjugated representation only in the case when this representation is defined by a pair $(0, l_1)$ or $(l_0, 0)$, that is, one from the two numbers l_0 and l_1 is equal to zero. We take $l_1 = 0$. In its turn, for the complex Clifford algebra $\mathbb{C}_n$ ($\overset{*}{\mathbb{C}}_n$), associated with the representation $\mathfrak{C}^{l_0+l_1-1,0}$ ($\mathfrak{C}^{0,l_0-l_1+1}$) of $\mathfrak{G}_+$, the equivalence of the representation to its conjugated representation induce a relation $\mathbb{C}_n = \overset{*}{\mathbb{C}}_n$, which is fulfilled only in the case when the algebra $\mathbb{C}_n = \mathbb{C} \otimes C\!\ell_{p,q}$ is reduced to its real subalgebra $C\!\ell_{p,q}$ ($p+q = n$). Thus, the restriction of the complex representation $\mathfrak{C}^{l_0+l_1-1,0}$ (or $\mathfrak{C}^{0,l_0-l_1+1}$) of the group $\mathbf{Spin}_+(1,3)$ onto the real representation, $(l_0, l_1) \to (l_0, 0)$, induce a restriction $\mathbb{C}_n \to C\!\ell_{p,q}$. Further, as is known, over the field $\mathbb{F} = \mathbb{R}$ at $p + q \equiv 0 \pmod 2$ there exist four types of real subalgebras $C\!\ell_{p,q}$: two types $p - q \equiv 0, 2 \pmod 8$ with the real division ring $\mathbb{K} \simeq \mathbb{R}$ and two types $p - q \equiv 4, 6 \pmod 8$ with the quaternionic division ring $\mathbb{K} \simeq \mathbb{H}$. Therefore, we have the following four classes of real representations of the group $\mathbf{Spin}_+(1,3)$:

$$\begin{aligned}
\mathfrak{R}_0^{l_0} &\leftrightarrow C\!\ell_{p,q},\ p - q \equiv 0 \pmod 8,\ \mathbb{K} \simeq \mathbb{R};\\
\mathfrak{R}_2^{l_0} &\leftrightarrow C\!\ell_{p,q},\ p - q \equiv 2 \pmod 8,\ \mathbb{K} \simeq \mathbb{R};\\
\mathfrak{H}_4^{l_0} &\leftrightarrow C\!\ell_{p,q},\ p - q \equiv 4 \pmod 8,\ \mathbb{K} \simeq \mathbb{H};\\
\mathfrak{H}_6^{l_0} &\leftrightarrow C\!\ell_{p,q},\ p - q \equiv 6 \pmod 8,\ \mathbb{K} \simeq \mathbb{H}.
\end{aligned} \tag{37}$$

We call the representations $\mathfrak{H}_4^{l_0}$ and $\mathfrak{H}_6^{l_0}$ *quaternionic representations* of the group $\mathbf{Spin}_+(1,3)$. On the other hand, over the field $\mathbb{F} = \mathbb{R}$ at $p + q \equiv 1 \pmod 2$ there exist four types of real subalgebras $C\!\ell_{p,q}$: two types $p - q \equiv 3, 7 \pmod 8$ with the complex division ring $\mathbb{K} \simeq \mathbb{C}$, one type $p - q \equiv 1 \pmod 8$ with the double division ring $\mathbb{K} \simeq \mathbb{R} \oplus \mathbb{R}$ and one type $p - q \equiv 5 \pmod 8$ with the double quaternionic division ring $\mathbb{K} \simeq \mathbb{H} \oplus \mathbb{H}$. Therefore, we have the following four classes of real representations of $\mathbf{Spin}_+(1,3)$:

$$\begin{aligned}
\mathfrak{C}_3^{l_0} &\leftrightarrow C\!\ell_{p,q},\ p - q \equiv 3 \pmod 8,\ \mathbb{K} \simeq \mathbb{C};\\
\mathfrak{C}_7^{l_0} &\leftrightarrow C\!\ell_{p,q},\ p - q \equiv 7 \pmod 8,\ \mathbb{K} \simeq \mathbb{C};\\
\mathfrak{R}_{0,2}^{l_0} \cup \mathfrak{R}_{0,2}^{l_0} &\leftrightarrow C\!\ell_{p,q},\ p - q \equiv 1 \pmod 8,\ \mathbb{K} \simeq \mathbb{R} \oplus \mathbb{R};\\
\mathfrak{H}_{4,6}^{l_0} \cup \mathfrak{H}_{4,6}^{l_0} &\leftrightarrow C\!\ell_{p,q},\ p - q \equiv 5 \pmod 8,\ \mathbb{K} \simeq \mathbb{H} \oplus \mathbb{H}.
\end{aligned} \tag{38}$$

Here $\mathfrak{R}^{l_0}_{0,2} \cup \mathfrak{R}^{l_0}_{0,2} \simeq \mathfrak{R}^{l_0}_{1}$ means that in virtue of an isomorphism $C\!\ell_{p,q} \simeq C\!\ell_{p,q-1} \oplus C\!\ell_{p,q-1}$ (or $C\!\ell_{p,q} \simeq C\!\ell_{q,p-1} \oplus C\!\ell_{q,p-1}$), where $p - q \equiv 1 \pmod 8$ and algebras $C\!\ell_{p,q-1}$ (or $C\!\ell_{q,p-1}$) are of the type $p - q \equiv 0 \pmod 8$ (or $p - q \equiv 2 \pmod 8$), a representation $\mathfrak{R}^{l_0}_{1}$ is equivalent to $\mathfrak{R}^{l_0}_{0} \cup \mathfrak{R}^{l_0}_{0}$ (or $\mathfrak{R}^{l_0}_{2} \cup \mathfrak{R}^{l_0}_{2}$). Analogously, we have $\mathfrak{H}^{l_0}_{4,6} \cup \mathfrak{H}^{l_0}_{4,6} \simeq \mathfrak{H}^{l_0}_{5}$ for representations of $\mathbf{Spin}_+(1,3)$ with the double quaternionic division ring $\mathbb{K} \simeq \mathbb{H} \oplus \mathbb{H}$. Therefore, all the real representations of $\mathbf{Spin}_+(1,3)$ are divided into the two sets: $\mathfrak{M}^+$ ($p - q \equiv 0, 2, 4, 6 \pmod 8$) and $\mathfrak{M}^-$ ($p - q \equiv 1, 3, 5, 7 \pmod 8$) which form a full system $\mathfrak{M} = \mathfrak{M}^+ \oplus \mathfrak{M}^-$ of the real representations of $\mathbf{Spin}_+(1,3)$. Let us find a relation of the number l_0 with the dimension of the real algebra $C\!\ell_{p,q}$. Using Karoubi Theorem, we obtain for the algebra $C\!\ell_{p,q}$ ($p + q \equiv 0 \pmod 2$) the following factorization:

$$C\!\ell_{p,q} \simeq \underbrace{C\!\ell_{s_i,t_j} \otimes C\!\ell_{s_i,t_j} \otimes \cdots \otimes C\!\ell_{s_i,t_j}}_{r \text{ times}}, \tag{39}$$

where $s_i, t_j \in \{0, 1, 2\}$. The algebra (39) corresponds to the representations $\mathfrak{R}^{l_0}_{0}$, $\mathfrak{R}^{l_0}_{2}$, $\mathfrak{H}^{l_0}_{4}$, $\mathfrak{H}^{l_0}_{6}$ of $\mathbf{Spin}_+(1,3)$. It is obvious that $l_0 = r/2$ and $n = 2r = p + q = 4l_0$, therefore, $l_0 = (p + q)/4$. When $p + q \equiv 1 \pmod 2$ we obtain

$$C\!\ell_{p,q} \simeq \underbrace{C\!\ell_{s_i,t_j} \otimes C\!\ell_{s_i,t_j} \otimes \cdots \otimes C\!\ell_{s_i,t_j}}_{r \text{ times}} \bigcup \underbrace{C\!\ell_{s_i,t_j} \otimes C\!\ell_{s_i,t_j} \otimes \cdots \otimes C\!\ell_{s_i,t_j}}_{r \text{ times}}. \tag{40}$$

In its turn, the algebra (40) is associated with the double representations $\mathfrak{R}^{l_0}_{0} \cup \mathfrak{R}^{l_0}_{0}$, $\mathfrak{R}^{l_0}_{2} \cup \mathfrak{R}^{l_0}_{2}$ and $\mathfrak{H}^{l_0}_{4} \cup \mathfrak{H}^{l_0}_{4}$, $\mathfrak{H}^{l_0}_{6} \cup \mathfrak{H}^{l_0}_{6}$ of the group $\mathbf{Spin}_+(1,3)$. □

5 Modulo 2 and 8 periodicities in particle physics

As is known, the Atiyah-Bott-Shapiro periodicity is based on the $\mathbb{Z}_2$-graded structure of the Clifford algebras $C\!\ell$. $\mathbb{Z}_2$-graded Clifford algebras and Brauer-Wall supergroups, considered in the section 2, allow one to define some cyclic relations on the representation system of $\mathfrak{G}_+ \simeq \mathbf{Spin}_+(1,3)$, which lead to interesting applications in particle physics (in some sense like to the Gell-Mann–Ne'eman eightfold way [37]).

1) Complex representations.

Graded central simple Clifford algebras over the field $\mathbb{F} = \mathbb{C}$ form two similarity classes, which, as it is easy to see, coincide with the two types of $\mathbb{C}_n$: $n \equiv 0, 1 \pmod 2$. The set of these two types (classes) forms a Brauer-Wall supergroup $BW_{\mathbb{C}} = G(\mathbb{C}_n, \gamma, \bigcirc)$ [17, 18], an action of which is formally equivalent to the action of the cyclic group $\mathbb{Z}_2$ (see section 2). Returning to the representations of $\mathfrak{G}_+$ (Theorem 3), we see that in virtue of correspondences $\mathbb{C}_n \leftrightarrow \mathfrak{C}$ ($n \equiv 0 \pmod 2$) and $\mathbb{C}_n \leftrightarrow \mathfrak{C} \cup \mathfrak{C}$ ($n \equiv 1 \pmod 2$) all the complex representations of the group $\mathfrak{G}_+$ form the full system $\mathfrak{M} = \mathfrak{M}^0 \oplus \mathfrak{M}^1$, where $\mathfrak{M}^0$ ($n \equiv 0 \pmod 2$) and $\mathfrak{M}^1$ ($n \equiv 1 \pmod 2$). The group action of $BW_{\mathbb{C}}$ can be transferred onto the system $\mathfrak{M} = \mathfrak{M}^0 \oplus \mathfrak{M}^1$. Indeed, a cyclic structure of the supergroup $BW_{\mathbb{C}}$ is defined by the transition $\mathbb{C}^+_n \xrightarrow{h} \mathbb{C}_n$, where the type of $\mathbb{C}_n$ is defined by the formula $n = h + 2r$, here $h \in \{0, 1\}$, $r \in \mathbb{Z}$ [15, 16]. Therefore, the action of $BW_{\mathbb{C}}$ on the system $\mathfrak{M}$ is defined by a transition $\overset{+}{\mathfrak{C}} \xrightarrow{h} \mathfrak{C}$, where $\overset{+}{\mathfrak{C}}{}^{l_0+l_1-1,0} \simeq \mathfrak{C}^{l_0+l_1-2,0}$ when $\mathfrak{C} \in \mathfrak{M}^0$ ($h = 1$) and $\overset{+}{\mathfrak{C}} = \left(\mathfrak{C}^{l_0+l_1-1,0} \cup \mathfrak{C}^{l_0+l_1-1,0}\right)^+ \sim \mathfrak{C}^{l_0+l_1-1,0}$ when $\mathfrak{C} \in \mathfrak{M}^1$ ($h = 0$), $\dim \mathfrak{C} = h + 2r$ ($\dim \mathfrak{C} = l_0 + l_1 - 1$ if $\mathfrak{C} \in \mathfrak{M}^0$ and $\dim \mathfrak{C} = 2(l_0 + l_1 - 1)$ if $\mathfrak{C} \in \mathfrak{M}^1$). For example, in virtue of the isomorphism $\mathbb{C}^+_2 \simeq \mathbb{C}_1$ the transition $\mathbb{C}^+_2 \to \mathbb{C}_2$ ($\mathbb{C}_1 \to \mathbb{C}_2$) induces on the system $\mathfrak{M}$ a transition $\overset{+}{\mathfrak{C}}{}^{1,0} \to \mathfrak{C}^{1,0}$ that in virtue of the isomorphism $\overset{+}{\mathfrak{C}}{}^{1,0} \simeq \mathfrak{C}^{0,0}$ is equivalent to $\mathfrak{C}^{0,0} \to \mathfrak{C}^{1,0}$ ($\mathfrak{C}^{0,0}$ is the one-dimensional representation of the group $\mathfrak{G}_+$) and, therefore, $h = 1$. In its turn, the transition



Fig. 5: An action of the supergroup $BW_{\mathbb{C}} = G(\mathbb{C}_n, \gamma, \bigcirc)$ on the system $\mathfrak{M} = \mathfrak{M}^0 \oplus \mathfrak{M}^1$ of complex representations $\mathfrak{C}$ of $\mathfrak{G}_+ \simeq \mathbf{Spin}_+(1,3)$.

$\mathbb{C}^+_3 \to \mathbb{C}_3$ ($\mathbb{C}_2 \to \mathbb{C}_3$) induces on the system $\mathfrak{M}$ a transition $(\mathfrak{C}^{1,0} \cup \mathfrak{C}^{1,0})^+ \to \mathfrak{C}^{1,0} \cup \mathfrak{C}^{1,0}$ and, therefore, $h = 0$. Thus, we see that a cyclic structure of the supergroup $BW_{\mathbb{C}}$ induces modulo 2 periodic relations on the system $\mathfrak{M}$, which can be explicitly shown on the Trautman diagram (see Fig. 5). Further, in virtue of the isomorphism $\mathbb{C}^+_4 \simeq \mathbb{C}_3$ the transition $\mathbb{C}^+_4 \to \mathbb{C}_4$ ($\mathbb{C}_3 \to \mathbb{C}_4$) induces on the system $\mathfrak{M}$ the following relation: $\overset{+}{\mathfrak{C}}{}^{2,0} \to \mathfrak{C}^{2,0}$ or $\mathfrak{C}^{1,0} \to \mathfrak{C}^{2,0}$ and, therefore, $h = 1$. Thus, a full rotation (cycle), presented on the diagram Fig. 5, corresponds to a transition from the representation $\mathfrak{C}^{1,0}$ of the spin 1/2, associated with the algebra $\mathbb{C}_2$, to the representation $\mathfrak{C}^{2,0}$ of the spin 1, associated with the algebra $\mathbb{C}_4$. It is easy to see that a sequence of transitions $\mathbb{C}^+_5 \to \mathbb{C}_5$ ($\mathbb{C}_4 \to \mathbb{C}_5$) and $\mathbb{C}^+_6 \to \mathbb{C}_6$ ($\mathbb{C}_5 \to \mathbb{C}_6$) leads to a transition from the representation $\mathfrak{C}^{2,0}$ of the spin 1 to the representation $\mathfrak{C}^{3,0}$ of the spin 3/2, associated with the algebra $\mathbb{C}_6$, and so on. Thus, the action of the supergroup $BW_{\mathbb{C}} = G(\mathbb{C}_n, \gamma, \bigcirc)$ on the system $\mathfrak{M} = \mathfrak{M}^0 \oplus \mathfrak{M}^1$ converts fermionic representations into bosonic representations and vice versa, that is, this action is equivalent to a supersymmetric action.

It is obvious that a cyclic structure over the algebras $\mathbb{C}_n$, defined by the supergroup $BW_{\mathbb{C}}$, is related immediately with a modulo 2 periodicity of complex Clifford algebras [14, 22]: $\mathbb{C}_{n+2} \simeq \mathbb{C}_n \otimes \mathbb{C}_2$. Therefore, we have the following relations for the complex representations $\mathfrak{C}$ and $\overset{*}{\mathfrak{C}}$:

$$\begin{aligned} \mathfrak{C}^{l_0+l_1,0} &\simeq \mathfrak{C}^{l_0+l_1-1,0} \otimes \mathfrak{C}^{1,0}, \\ \mathfrak{C}^{0,l_0-l_1} &\simeq \mathfrak{C}^{0,l_0-l_1+1} \otimes \mathfrak{C}^{0,-1}. \end{aligned}$$

Correspondingly, for the representation $\mathfrak{C} \otimes \overset{*}{\mathfrak{C}}$ we obtain

$$\mathfrak{C}^{l_0+l_1,l_0-l_1} \simeq \mathfrak{C}^{l_0+l_1-1,l_0-l_1+1} \otimes \mathfrak{C}^{1,-1}.$$

Thus, the action of the supergroup $BW_{\mathbb{C}}$ forms a cycle of the period 2 on the system $\mathfrak{M}$, where a basic cycle generating factor is the fundamental representation $\mathfrak{C}^{1,0}$ ($\mathfrak{C}^{0,-1}$) of the group $\mathfrak{G}_+$.

2) Real representations.

A restriction of the complex representations $\mathfrak{C}$ onto the real representations $\mathfrak{R}$ and $\mathfrak{H}$ leads to a much more high-graded periodicity over the field $\mathbb{F} = \mathbb{R}$. Physical fields, defined within such representations, describe neutral particles, or particles occurring in the rest such as atomic nuclei. So, the Clifford algebra $\mathcal{C}\ell_{p,q}$ is central simple if $p - q \not\equiv 1, 5 \pmod 8$. Graded central simple Clifford algebras over the field $\mathbb{F} = \mathbb{R}$ form eight similarity classes, which, as it is easy to see, coincide with the eight types of the algebras $\mathcal{C}\ell_{p,q}$. The set of these eight types (classes) forms the Brauer-Wall supergroup $BW_{\mathbb{R}} = G(\mathcal{C}\ell_{p,q}, \gamma, \bigcirc)$ [17, 18], a cyclic structure of which is formally equivalent to the action of cyclic group $\mathbb{Z}_8$ (see section 2). Therefore, in virtue



Fig. 6: An action of the supergroup $BW_{\mathbb{R}} = G(C\!\ell_{p,q},\gamma,\circ)$ on the system $\mathfrak{M} = \mathfrak{M}^+\oplus\mathfrak{M}^-$ of real representations $\mathfrak{D}$ of $\mathfrak{G}_+\simeq$ $\mathbf{Spin}_+(1,3)$, $l_0=\frac{p+q}{4}$.

of the correspondences (37) and (38) an action of $BW_{\mathbb{R}}$ can be transferred onto the system $\mathfrak{M} = \mathfrak{M}^+\oplus\mathfrak{M}^-$. In its turn, a cyclic structure of the supergroup $BW_{\mathbb{R}}$ is defined by a transition $C\!\ell^+_{p,q} \xrightarrow{h} C\!\ell_{p,q}$, where the type of $C\!\ell_{p,q}$ is defined by the formula $q-p=h+8r$, here $h\in\{0,\ldots,7\}$, $r\in\mathbb{Z}$ [15, 16]. Thus, the action of $BW_{\mathbb{R}}$ on the system $\mathfrak{M}$ is defined by a transition $\overset{+}{\mathfrak{D}}{}^{l_0} \xrightarrow{h} \mathfrak{D}^{l_0}$, where $\mathfrak{D}^{l_0} = \left\{\mathfrak{R}^{l_0}_{0,2}, \mathfrak{H}^{l_0}_{4,6}, \mathfrak{C}^{l_0}_{3,7}, \mathfrak{R}^{l_0}_{0,2}\cup\mathfrak{R}^{l_0}_{0,2}, \mathfrak{H}^{l_0}_{4,6}\cup\mathfrak{H}^{l_0}_{4,6}\right\}$, and $\overset{+}{\mathfrak{D}}{}^{r/2}\simeq\mathfrak{D}^{\frac{r-1}{2}}$ if $\mathfrak{D}\in\mathfrak{M}^+$ and $\left(\mathfrak{D}^{r/2}\cup\mathfrak{D}^{r/2}\right)^+\simeq\mathfrak{D}^{r/2}$ if $\mathfrak{D}\in\mathfrak{M}^-$. Therefore, the cyclic structure of the supergroup $BW_{\mathbb{R}} = G(C\!\ell_{p,q},\gamma,\circ)$ induces on the system $\mathfrak{M}$ modulo 8 periodic relations, which can be shown explicitly on the diagram Fig. 6. Let us consider in detail one action cycle of the supergroup $BW_{\mathbb{R}}$ on the system $\mathfrak{M} = \mathfrak{M}^+\oplus\mathfrak{M}^-$. In virtue of an isomorphism $C\!\ell^+_{1,1}\simeq C\!\ell_{1,0}$ a transition $C\!\ell^+_{1,1}\rightarrow C\!\ell_{1,1}$ ($C\!\ell_{1,0}\rightarrow C\!\ell_{1,1}$) induces on the system $\mathfrak{M}$ a transition $\mathfrak{R}^0_0\cup\mathfrak{R}^0_0\rightarrow\mathfrak{R}^{\frac{1}{2}}_0$, where $C\!\ell_{1,0}\simeq C\!\ell_{0,0}\cup C\!\ell_{0,0}$ is an algebra of double numbers, $\mathfrak{R}^0_0$ is the field of real numbers, $\mathfrak{R}^{\frac{1}{2}}_0$ is a real representation of the spin 1/2, associated with the algebra $C\!\ell_{1,1}$. At this point, $h=0$ and $r=0$ (an original point of the first cycle). Further, in virtue of $C\!\ell^+_{1,2}\simeq C\!\ell_{1,1}$ a transition $C\!\ell^+_{1,2}\rightarrow C\!\ell_{1,2}$ ($C\!\ell_{1,1}\rightarrow C\!\ell_{1,2}$) induces on the system $\mathfrak{M}$ a transition $\mathfrak{R}^{\frac{1}{2}}_0\rightarrow\mathfrak{R}^{\frac{1}{2}}_0\cup\mathfrak{R}^{\frac{1}{2}}_0$, at this point, $h=1$ and $r=0$. The following transition $C\!\ell^+_{1,3}\rightarrow C\!\ell_{1,3}$ ($C\!\ell_{1,2}\rightarrow C\!\ell_{1,3}$) in virtue of $C\!\ell^+_{1,3}\simeq C\!\ell_{1,2}$ leads to a transition $\mathfrak{R}^{\frac{1}{2}}_0\cup\mathfrak{R}^{\frac{1}{2}}_0\rightarrow\mathfrak{H}^1_6$, where $\mathfrak{H}^1_6$ is a quaternionic representation of the spin 1, associated with the algebra $C\!\ell_{1,3}$. At this transition we have $h=2$ and $r=0$. In virtue of an isomorphism $C\!\ell^+_{1,4}\simeq C\!\ell_{1,3}$ a transition $C\!\ell^+_{1,4}\rightarrow C\!\ell_{1,4}$ ($C\!\ell_{1,3}\rightarrow C\!\ell_{1,4}$) induces on the system $\mathfrak{M}$ a transition $\mathfrak{H}^1_6\rightarrow\mathfrak{H}^1_6\cup\mathfrak{H}^1_6$, at this point, $h=3$ and $r=0$. Further, in virtue of $C\!\ell^+_{1,5}\simeq C\!\ell_{1,4}$ a transition $C\!\ell^+_{1,5}\rightarrow C\!\ell_{1,5}$ ($C\!\ell_{1,4}\rightarrow C\!\ell_{1,5}$) induces on the system $\mathfrak{M}$ a transition $\mathfrak{H}^1_6\cup\mathfrak{H}^1_6\rightarrow\mathfrak{H}^{\frac{3}{2}}_4$, where $\mathfrak{H}^{\frac{3}{2}}_4$ is a quaternionic representation of the spin 3/2, associated with the algebra $C\!\ell_{1,5}$. At this transition we have $h=4$ and $r=0$. The following transition $C\!\ell^+_{1,6}\rightarrow C\!\ell_{1,6}$ ($C\!\ell_{1,5}\rightarrow C\!\ell_{1,6}$) in virtue of $C\!\ell^+_{1,6}\simeq C\!\ell_{1,5}$ leads to a transition $\mathfrak{H}^{\frac{3}{2}}_4\rightarrow\mathfrak{H}^{\frac{3}{2}}_4\cup\mathfrak{H}^{\frac{3}{2}}_4$, at this point, $h=5$ and $r=0$. The transition $C\!\ell^+_{1,7}\rightarrow C\!\ell_{1,7}$ ($C\!\ell_{1,6}\rightarrow C\!\ell_{1,7}$) in virtue of $C\!\ell^+_{1,7}\simeq C\!\ell_{1,6}$ induces on the system $\mathfrak{M}$ a transition $\mathfrak{H}^{\frac{3}{2}}_4\cup\mathfrak{H}^{\frac{3}{2}}_4\rightarrow\mathfrak{R}^2_2$, where $\mathfrak{R}^2_2$ is a real representation of the spin 2, associated with the algebra $C\!\ell_{1,7}$. At this transition we have $h=6$ and $r=0$. Finally, a transition $C\!\ell^+_{1,8}\rightarrow C\!\ell_{1,8}$ ($C\!\ell_{1,7}\rightarrow C\!\ell_{1,8}$) finishes the first cycle ($h=7$, $r=0$) and in virtue of $C\!\ell^+_{1,8}\simeq C\!\ell_{1,7}$ induces on the system $\mathfrak{M}$ the following transition: $\mathfrak{R}^2_2\rightarrow\mathfrak{R}^2_2\cup\mathfrak{R}^2_2$. It is obvious that a new cycle ($h=0$, $r=1$) is started with a transition $\mathcal{C}\ell^+_{1,9}\rightarrow\mathcal{C}\ell_{1,9}$ ($\mathcal{C}\ell_{1,8}\rightarrow\mathcal{C}\ell_{1,9}$) and so on.

Further, a cyclic structure over the algebras $\mathcal{C}\ell_{p,q}$, defined by the supergroup $BW_{\mathbb{R}}$, is related immediately with the Atiyah-Bott-Shapiro periodicity [14]. In accordance with [14] the Clifford algebra over the field $\mathbb{F}=\mathbb{R}$ is modulo 8 periodic: $\mathcal{C}\ell_{p+8,q}\simeq\mathcal{C}\ell_{p,q}\otimes\mathcal{C}\ell_{8,0}$ ($\mathcal{C}\ell_{p,q+8}\simeq\mathcal{C}\ell_{p,q}\otimes\mathcal{C}\ell_{0,8}$). Therefore, the following relation takes place:

$$\mathfrak{D}^{l_0+2}\simeq\mathfrak{D}^{l_0}\otimes\mathfrak{R}^2_0,$$

since $\mathfrak{R}^2_0\leftrightarrow\mathcal{C}\ell_{8,0}$ ($\mathcal{C}\ell_{0,8}$). In virtue of the theorem 2 from (39) it follows that $\mathcal{C}\ell_{8,0}\simeq\mathcal{C}\ell_{2,0}\otimes\mathcal{C}\ell_{0,2}\otimes\mathcal{C}\ell_{0,2}\otimes\mathcal{C}\ell_{2,0}$ ($\mathcal{C}\ell_{0,8}\simeq\mathcal{C}\ell_{0,2}\otimes\mathcal{C}\ell_{2,0}\otimes\mathcal{C}\ell_{2,0}\otimes\mathcal{C}\ell_{0,2}$). A minimal left ideal of the algebra $\mathcal{C}\ell_{8,0}$ is the spinspace $\mathbb{S}_{16}$, which in virtue of the real ring $\mathbb{K}\simeq\mathbb{R}$ is defined within a full matrix algebra $\mathrm{Mat}_{16}(\mathbb{R})$. At first glance, from the factorization of the algebra $\mathcal{C}\ell_{8,0}$ it follows that $\mathrm{Mat}_2(\mathbb{R})\otimes\mathbb{H}\otimes\mathbb{H}\otimes\mathrm{Mat}_2(\mathbb{R})\not\simeq\mathrm{Mat}_{16}(\mathbb{R})$, but it is wrong, since there exists an isomorphism $\mathbb{H}\otimes\mathbb{H}\simeq\mathrm{Mat}_4(\mathbb{R})$ (see Appendix B in [38]). On the other hand, in terms of minimal left ideals the modulo 8 periodicity is defined as follows:

$$\mathbb{S}_{n+8}\simeq\mathbb{S}_n\otimes\mathbb{S}_{16}.$$

In virtue of the mapping $\gamma_{8,0}:\mathcal{C}\ell_{8,0}\rightarrow\mathrm{Mat}_2(\mathbb{O})$ [39, 40], the latter relation can be written in the form

$$\mathbb{S}_{n+8}\simeq\mathbb{S}_n\otimes\mathbb{O}^2,$$

where $\mathbb{O}$ is *an octonion algebra*. Because the algebra $\mathcal{C}\ell_{8,0}\simeq\mathcal{C}\ell_{0,8}$ admits an octonionic representation, then in virtue of the modulo 8 periodicity octonionic representations can be defined for all high dimensions and, therefore, on the system $\mathfrak{M}=\mathfrak{M}^+\oplus\mathfrak{M}^-$ of real representations of the group $\mathfrak{G}_+$ we have a relation

$$\mathfrak{D}^{l_0+2}\simeq\mathfrak{D}^{l_0}\otimes\mathfrak{O},$$

where $\mathfrak{O}$ is *an octonionic representation* of the group $\mathfrak{G}_+=\mathbf{Spin}_+(1,3)$ ($\mathfrak{O}\sim\mathfrak{R}^0_4$).

Acknowledgements

I am deeply grateful to Prof. C. T. C. Wall and Prof. A. Trautman for sending me their interesting works.

References

[1] W. K. Clifford, Applications of Grassmann's extensive algebra, Amer. J. Math. **1** (1878) 350–358.

[2] W. R. Hamilton, Lectures on Quaternions, Hodges and Smith, Dublin, 1853.

[3] H. Grassmann, Die Ausdehnungslehre, Enslin, Berlin, 1862.

[4] W. K. Clifford, On the classification of geometric algebras, in R. Tucker, ed., Mathematical Papers by William Kingdon Clifford, Macmillan, London, 1982, pp. 397-401.

[5] R. Lipschitz, Untersuchungen über die Summen von Quadraten, Max Cohen und Sohn, Bonn, 1886.

[6] E. Cartan, Nombres complexes, in J. Molk, ed. : Encyclopédie des sciences mathématiques, Tome I, Vol.1, Fasc 4, art. 15, (1908) pp. 329-468.

[7] E. Witt, Theorie der quadratischen Formen in beliebigen Körpern, J. Reine Angew. Math. **176** (1937) 31–44.

[8] C. Chevalley, The Algebraic Theory of Spinors, Columbia University Press, New York, 1954.

[9] A. Crumeyrolle, Orthogonal and Symplectic Clifford Algebras, Spinor Structures, Kluwer Acad. Publ., Dordrecht, 1991.

[10] I. R. Porteous, Clifford Algebras and Classical Groups, Cambridge University Press, Cambridge, 1995.

[11] P. Lounesto, Clifford Algebras and Spinors, Cambridge University Press, Cambridge, 1997.

[12] S. Bochner, Formal Lie Groups, Ann. Math. **47** (1946) 192–212.

[13] F. A. Berezin, G. I. Kac, Lie groups with commuting and anticommuting parameters, Mat. Zbornik **82** (1970) 343–359.

[14] M. F. Atiyah, R. Bott, A. Shapiro, Clifford modules, Topology **3**, Suppl. 1, (1964) 3–38.

[15] P. Budinich, A. Trautman, An introduction to the spinorial chessboard, J. Geom. Phys. **4** (1987) 363–390.

[16] P. Budinich, A. Trautman, The Spinorial Chessboard, Springer, Berlin, 1988.

[17] C. T. C. Wall, Graded Brauer Groups, J. Reine Angew. Math. **213** (1964) 187–199.

[18] P. Lounesto, Scalar Products of Spinors and an Extension of Brauer-Wall Groups, Found. Phys. **11** (1981) 721–740.

[19] D. M. Gitman, A. L. Shelepin, Field on the Poincaré group and quantum description of orientable objects, Eur. Phys. J. C. **61** (2009) 111–139; arXiv:0901.2537 [hep-th] (2009).

[20] D. M. Gitman, A. L. Shelepin, Classification of quantum relativistic orientable objects, Phys. Scr. **83** (2011) 015103; arXiv: 1001.5290 [hep-th] (2010).

[21] C. Chevalley, The construction and study of certain important algebras, Publications of Mathematical Society of Japan No 1, Herald Printing, Tokyo, 1955.

[22] M. Karoubi, K-Theory. An Introduction, Springer-Verlag, Berlin, 1979.

[23] P. K. Rashevskii, The Theory of Spinors, (in Russian) Uspekhi Mat. Nauk **10** (1955), 3–110; English translation in Amer. Math. Soc. Transl. (Ser. 2) **6** (1957) 1.

[24] I. R. Porteous, Topological Geometry, van Nostrand, London, 1969.

[25] I. M. Gel'fand, R. A. Minlos, Z. Ya. Shapiro, Representations of the Rotation and Lorentz Groups and their Applications, Pergamon Press, Oxford, 1963.

[26] B. L. van der Waerden, Die Gruppentheoretische Methode in der Quantenmechanik, Springer, Berlin, 1932.

[27] I. M. Gel'fand, A. M. Yaglom, General relativistic–invariant equations and infinite–dimensional representations of the Lorentz group, Zh. Ehksp. Teor. Fiz. **18** (1948) 703–733.

[28] V. V. Varlamov, General Solutions of Relativistic Wave Equations, Int. J. Theor. Phys. **42** (2003) 583–633; arXiv:math-ph/0209036 (2002).

[29] V. V. Varlamov, Relativistic wavefunctions on the Poincare group, J. Phys. A: Math. Gen. **37** (2004) 5467–5476; arXiv:math-ph/0308038 (2003).

[30] V. V. Varlamov, Maxwell field on the Poincaré group, Int. J. Mod. Phys. A. **20** (2005) 4095–4112; arXiv:math-ph/0310051 (2003).

[31] M. Fierz, W. Pauli, On Relativistic Wave Equations of Particles of Arbitrary Spin in an Electromagnetic Field. Proc. Roy. Soc. (London) A. **173** (1939) 211–232.

[32] V. V. Varlamov, General Solutions of Relativistic Wave Equations II: Arbitrary Spin Chains, Int. J. Theor. Phys. **46** (2007) 741–805; arXiv:math-ph/0503058 (2005).

[33] M. A. Naimark, Linear Representations of the Lorentz Group, Pergamon, London, 1964.

[34] Yu. B. Rumer, A. I. Fet, Group Theory and Quantized Fields, Nauka, Moscow, 1977 [in Russian].

[35] L. de Broglie, Theorie Generale des Particules a Spin (Methode de Fusion), Gauthier-Villars, Paris, 1943.

[36] E. P. Wigner, Unitary Representations of the Inhomogeneous Lorentz Group Including Reflections in Group Theoretical Concepts and Methods in Elementary Particle Physics, Ed. F. Gürsey, Gordon & Breach, New York, 1964.

[37] M. Gell-Mann, Y. Ne'eman, The Eightfold Way, Benjamin, New York, 1964.

[38] M. Berg, C. DeWitt-Morette, S. Gwo, E. Kramer, The Pin Groups in Physics: C, P, and T, Rev. Math. Phys. **13** (2001) 953–1034; arXiv:math-ph/0012006 (2000).

[39] C. A. Manogue, J. Schray, Octonionic representations of Clifford algebras and triality, Found. Phys. **26** (1996) 17–70; arXiv:hep-th/9407179 (1994).

[40] J. C. Baez, The Octonions, Bull. Am. Math. Soc. **39** (2002) 145–205; arXiv:math.RA/0105155 (2001).